arXiv:1110.6653v1 [math.AC] 30 Oct 2011

Betti numbers of the path ideals of cycles and lines

Ali Alilooee[*] Sara Faridi[†]

November 1, 2011

1 Introduction

Path ideals of graphs were first introduced by Conca and De Negri [3] in the context of monomial ideals of linear type. Simply put, path ideals are ideals whose monomial generators correspond to vertices in paths of a given length in a graph. Conca and De Negri showed that path ideals of trees have normal and Cohen-Macaulay Rees rings. More recently Bouchat, Hà and O'Keefe [1] and He and Van Tuyl [7] studied invariants related to resolutions of path ideals of certain graphs.

In his thesis, Jacques [8] used beautiful techniques to compute Betti numbers of edge ideals of several classes of graphs. Edge ideals can be considered as path ideals of length 2.

Our paper extends Jacque's techniques to higher dimensions. Essentially, we consider the path ideal of a graph as a disjoint union of connected components. We then use homological methods to glue these components back together, and using Hochster's formula we compute all graded Betti numbers, projective dimension and regularity of path ideals of cycles and lines. Our paper extends the results of Bouchat, Hà and O'Keefe [1], where they compute some special cases of these Betti numbers, and gives a new proof (as a corollary of our main theorem) of the formula for the projective dimension in the case of lines that appears in He and Van Tuyl [7].

The paper is organized as follows. In Section 2 we recall some notation and basic algebraic and combinatorial concepts used in other next chapters. In Section 3 we study the connected components of path ideals which will provide us with the key to our homological computations later in Section 4. Section 5 is where we apply the homological results of Section 4 along with some combinatorics to compute all the Betti numbers and projective dimension of path ideals of lines and cycles.

While working on this paper the computer algebra systems CoCoA [11] and Macaulay2 [4] were used to test examples. We acknowledge the immense help that they have provided us in this project.

2 Preliminaries

Throughout we assume that K is a field and $R = K\left[x_1, \ldots, x_n\right]$ is a polynomial ring in n variables.

Simplicial complexes and monomial ideals

Definition 2.1. An abstract **simplicial complex** on vertex set $\mathcal{X} = \{x_1, \ldots, x_n\}$ is a collection Δ of subsets of $\mathcal{X}$ satisfying

---

[*] Department of Mathematics and Statistics, Dalhousie University, Halifax, Canada, alilooee@mathstat.dal.ca.

[†] Department of Mathematics and Statistics, Dalhousie University, Halifax, Canada, faridi@mathstat.dal.ca.

i. $\{x_i\} \in \Delta$ for all i,

ii. $F \in \Delta, G \subset F \Longrightarrow G \in \Delta$.

The elements of Δ are called **faces** of Δ and the maximal faces under inclusion are called **facets** of Δ. We denote the simplicial complex Δ with facets $F_1, \ldots, F_s$ by $\langle F_1, \ldots, F_s \rangle$. We call $\{F_1, \ldots, F_s\}$ the facet set of Δ and is denoted by $F(\Delta)$. The vertex set of Δ is denoted by Vert(Δ).

Definition 2.2. A **subcollection** of a simplicial complex Δ with vertex set $\mathcal{X}$ is a simplicial complex whose facet set is a subset of the facet set of Δ. For $\mathcal{Y} \subseteq \mathcal{X}$, an **induced subcollection** of Δ on $\mathcal{Y}$, denoted by $\Delta_{\mathcal{Y}}$, is the simplicial complex whose vertex set is a subset of $\mathcal{Y}$ and facet set is

$$\{F \in F(\Delta) \mid F \subseteq \mathcal{Y}\}.$$

If F is a face of $\Delta = \langle F_1, \ldots, F_s \rangle$, we define the **complement** of F in Δ to be

$$F^c_{\mathcal{X}} = \mathcal{X} \setminus F \quad \text{and} \quad \Delta^c_{\mathcal{X}} = \langle (F_1)^c_{\mathcal{X}}, \ldots, (F_s)^c_{\mathcal{X}} \rangle.$$

Also if $\mathcal{X} \subsetneqq \text{Vert}(\Delta)$, then $\Delta^c_{\mathcal{X}} = (\Delta_{\mathcal{X}})^c_{\mathcal{X}}$.

Definition 2.3. Let $R = K[x_1, \ldots, x_n]$ be a polynomial ring over a field K, and I an ideal in R minimally generated by square-free monomials $m_1, \ldots, m_s$. One can associate two simplicial complexes to I.

i. The **Stanley-Reisner complex** Δ_I associated to I has vertex set $V = \{x_i \mid x_i \notin I\}$ and is defined as
$$\Delta_I = \{\{x_{i_1}, \ldots, x_{i_k}\} \mid i_1 < i_2 < \cdots < i_k, x_{i_1} \ldots x_{i_k} \notin I\}.$$

ii. The **facet complex** $\Delta(I)$ associated to I has vertex set $\{x_1, \ldots, x_n\}$ and is defined as
$$\Delta(I) = \langle F_1, \ldots, F_s \rangle \text{ where } F_i = \{x_j \mid x_j | m_i,\ 1 \leq j \leq n\},\ 1 \leq i \leq s.$$

Conversely to a simplicial complex Δ one can associate two monomial ideals.

Definition 2.4. Let Δ be a simplicial complex with vertex set $x_1, \ldots, x_n$ and $R = K[x_1, \ldots, x_n]$ be a polynomial ring over a field K.

i. The **Stanley-Reisner** ideal of Δ is defined as
$$I_\Delta = (\prod_{x \in F} x \mid F \notin \Delta).$$

ii. The **facet ideal** of Δ is defined as
$$I(\Delta) = (\prod_{x \in F} x \mid \ F \text{ is a facet of} \Delta).$$

Note that there is a one-to-one correspondence between monomial ideals and simplicial complexes via each of these methods.

Definition 2.5. Let Δ be a simplicial complex with vertex set $\mathcal{X}$. The **Alexander Dual** Δ^* is defined to be the simplicial complex with faces

$$\Delta^* = \{F \subset \mathcal{X} \mid F^c_{\mathcal{X}} \text{ is not face of } \Delta\}.$$

To prove some of the results in this paper we need the definition of a cone and its properties.

Definition 2.6. Let Δ_1 and Δ_2 be two simplicial complexes on disjoint vertex sets V and W. The **join** $\Delta_1 * \Delta_2$ is the simplicial complex on $V \bigsqcup W$ with faces $F \cup G$ where $F \in \Delta_1, G \in \Delta_2$. The **cone** $Cn(\Delta)$ of Δ is the simplicial complex $\omega * \Delta$, where ω is a new vertex.

Given a simplicial complex Δ, we denote by $\mathcal{C}.(\Delta)$ the reduced chain complex and by $\widetilde{H}_i(\Delta) = Z_i(\Delta)/B_i(\Delta)$ the i - th reduced homology groups of Δ with coefficients in field K. For the proof of the following fact, see for example Villarreal [12], Proposition 5.2.5.

Proposition 2.7. *If Δ is a simplicial complex we have $\widetilde{H}_i(Cn(\Delta)) = 0$ for all i.*

Betti numbers

For any homogeneous ideal I of the polynomial ring $R = K[x_1, \ldots, x_n]$ there exists a **graded minimal finite free resolution**

$$0 \to \bigoplus_d R(-d)^{\beta_{p,d}} \to \cdots \to \bigoplus_d R(-d)^{\beta_{1,d}} \to R \to R/I \to 0$$

of R/I in which $R(-d)$ denotes the graded free module obtained by shifting the degrees of elements in R by d. The numbers $\beta_{i,d}$, which we shall refer to as the i-th $\mathbb{N}$-**graded Betti numbers** of degree d of R/I, are independent of the choice of graded minimal finite free resolution.

For computing the $\mathbb{N}$-graded Betti numbers of the Stanley-Reisner ring of a simplicial complex we use an equivalent form of Hochster's formula.

Theorem 2.8. *Let $R = K[x_1, \ldots, x_n]$ be a polynomial ring over a field K, and I be a pure square-free monomial ideal in R. Then the $\mathbb{N}$-graded Betti numbers of R/I are given by*

$$\beta_{i,d}(R/I) = \sum_{\Gamma \subset \Delta(I), |Vert(\Gamma)| = d} \dim_K \widetilde{H}_{i-2}(\Gamma^c_{Vert(\Gamma)})$$

where the sum is taken over the induced subcollections Γ of $\Delta(I)$ which have d vertices.

Proof. Hochster's formula (see for example Corollary 5.12 of [9]) says

$$\beta_{i,d}(R/I) = \sum_{W \subset \mathrm{Vert}(\Delta_I), |W| = d} \dim_K \widetilde{H}_{d-i-1}(\{\Delta_I\}_W)$$

where $\{\Delta_I\}_W = \{F \in \Delta_I \mid F \subset W\}$. On the other hand from [2] Lemma 5.5.3 we have

$$\widetilde{H}_{d-i-1}(\{\Delta_I\}_W) \cong \widetilde{H}^{i-2}(\{\Delta_I\}_W{}^*) \cong \widetilde{H}_{i-2}(\{\Delta_I\}_W{}^*). \tag{2.1}$$

Suppose $m_1, m_2, \ldots, m_r$ is a minimal monomial generating set for I and correspondingly, $\Delta(I) = \langle F_1, \ldots, F_s \rangle$. We now claim $\{\Delta_I\}_W{}^* = (\Delta(I)_W)^c_W$ for $W \subset \mathrm{Vert}(\Delta_I)$.

$$\begin{aligned}
F \in \{\Delta_I\}_W{}^* &\iff W \setminus F \notin \{\Delta_I\}_W \\
&\iff W \backslash F \notin \Delta_I \\
&\iff \prod_{x \in W \backslash F} x \in I = (m_1, m_2, \ldots, m_r) \\
&\iff m_s | \prod_{x \in W \backslash F} x, \text{ for some } s \in \{1, \ldots, r\} \\
&\iff F_s \subset W \backslash F \subset W, \text{ for some } s \in \{1, \ldots, r\} \\
&\iff F \subset W \backslash F_s \in (\Delta(I)_W)^c_W, \text{ for some } s \in \{1, \ldots, r\}.
\end{aligned}$$

Now Hochster's formula and (2.1) imply that

$$\beta_{i,d}(R/I) = \sum_{W \subset \text{Vert}(\Delta), |W|=d} dim_K \widetilde{H}_{i-2}((\Delta(I)_W)^c_W).$$

If we assume Vert$(\Delta(I)_W) \neq W$ then clearly we have $(\Delta(I)_W)^c_W$ is a cone and by Proposition 2.7 it contributes zero to the sum. So we have

$$\beta_{i,d}(R/I) = \sum_{\Gamma \subset \Delta(I), |\text{Vert}(\Gamma)|=d} dim_K \widetilde{H}_{i-2}(\Gamma^c_{\text{Vert}(\Gamma)}).$$

where the sum is taken over the induced subcollections Γ of $\Delta(I)$ which have d vertices. $\square$

Based on Theorem 2.8, from here on all induced subcollections $\Gamma = \Delta_{\mathcal{Y}}$ of a simplicial complex Δ that we consider will have the property that $\mathcal{Y} = $ Vert(Γ).

3 Path ideals and runs

We now focus on path ideals, path complexes, and their structures.

Definition 3.1. Let $G = (\mathcal{X}, E)$ be a finite simple graph and t be an integer such that $t \geq 2$. If x and y are two vertices of G, a **path** of length $(t-1)$ from x to y is a sequence of vertices $x = x_{i_1}, \ldots, x_{i_t} = y$ of G such that $\{x_{i_j}, x_{i_{j+1}}\} \in E$ for all $j = 1, 2, \ldots, t-1$. We define the **path ideal** of G, denoted by $I_t(G)$ to be the ideal of $K[x_1, \ldots, x_n]$ generated by the monomials of the form $x_{i_1}x_{i_2}\ldots x_{i_t}$ where $x_{i_1}, x_{i_2}, \ldots, x_{i_t}$ is a path in G. The facet complex of $I_t(G)$, denoted by $\Delta_t(G)$, is called the **path complex** of the graph G.

Two special cases that we will be considering in this paper are when G is a **cycle** C_n, or a **line graph** L_n on vertices $\{x_1, \ldots, x_n\}$.

$$C_n = \langle x_1x_2, \ldots, x_{n-1}x_n, x_nx_1 \rangle \text{ and } L_n = \langle x_1x_2, \ldots, x_{n-1}x_n \rangle.$$

Example 3.2. Consider the cycle C_7 with vertex set $\mathcal{X} = \{x_1, \ldots, x_7\}$



Then we have

$$\begin{aligned} I_4(C_7) =& (x_1x_2x_3x_4, x_2x_3x_4x_5, x_3x_4x_5x_6, x_4x_5x_6x_7, x_1x_5x_6x_7, x_1x_2x_6x_7, x_1x_2x_3x_7) \\ \Delta_4(C_7) =& \langle \{x_1,x_2,x_3,x_4\}, \{x_2,x_3,x_4,x_5\}, \{x_3,x_4,x_5,x_6\}, \{x_4,x_5,x_6,x_7\}, \\ & \{x_1,x_5,x_6,x_7\}, \{x_1,x_2,x_6,x_7\}, \{x_1,x_2,x_3,x_7\} \rangle. \end{aligned}$$

Notation 3.3. Let i and n be two positive integers. For (a set of) labeled objects we use the notation $\mod n$ to denote

$$x_i \mod n = \{x_j \mid 1 \leq j \leq n, i \equiv j \mod n\}$$

and

$$\{x_{u_1}, x_{u_2}, \ldots, x_{u_t}\} \mod n = \{x_{u_j} \mod n \mid j = 1, 2, \ldots, n\}.$$

Note 3.4. Let C_n be a cycle on vertex set $\mathcal{X} = \{x_1, \dots, x_n\}$ and $t < n$. The facets of the path complex $\Delta_t(C_n) = \langle F_1, \dots, F_n \rangle$ can be labeled as

$$F_1 = \{x_1, \dots, x_t\}, \dots, F_{n-(t-1)} = \{x_{n-(t-1)}, \dots, x_n\}, \dots, F_n = \{x_1, \dots, x_{t-1}, x_n\}$$

such that $F_i = \{x_i, x_{i+1}, \dots, x_{i+t-1}\} \mod n$ for all $1 \leq i \leq n$. This labeling is called the **standard labeling** of $\Delta_t(C_n)$.

Since for each $1 \leq i \leq n$ we have

$$F_{i+1} \setminus F_i = \{x_{t+i}\} \quad \text{and} \quad F_i \setminus F_{i+1} = \{x_i\} \qquad \mod n,$$

it follows that $|F_i \setminus F_{i+1}| = 1$ and $|F_{i+1} \setminus F_i| = 1 \quad \mod n$ for all $1 \leq i \leq n-1$.

It is clear that each induced subgraph of a graph-cycle is a disjoint union of paths. Borrowing the terminology from S. Jacques [8], we call the path complex of a line a "run", and show that every induced subcollection of the path complex of a cycle is a disjoint union of runs.

Definition 3.5. Given an integer t, we define a **run** to be the path complex of a line graph. A run which has p facets is called a **run of length** p and corresponds to $\Delta_t(L_{p+t-1})$. Therefore a run of length p has $p + t - 1$ vertices.

Proposition 3.6 below shows that every proper induced subcollection of a path complex is a disjoint union of runs.

Proposition 3.6. *Let C_n be a cycle with vertex set $\mathcal{X} = \{x_1, \dots . x_n\}$ and $2 \leq t < n$. Let Γ be a proper induced connected subcollection of $\Delta_t(C_n)$ on $U \subsetneq \mathcal{X}$. Then Γ is of the form $\Delta_t(L_{|\Gamma|})$, where $L_{|\Gamma|}$ is the line graph on $|\Gamma|$ vertices.*

Proof. Suppose $\Delta_t(C_n) = \langle F_1, \dots, F_n \rangle$ has standard labeling and $\Gamma = \langle F_{i_1}, \dots, F_{i_r} \rangle$. It is clear that there exists the facet $F_a \in \Gamma$ for $1 \leq a \leq n$ such that $F_{a+1} \notin \Gamma \mod n$, because otherwise $\Gamma = \Delta_t(C_n)$. Therefore from Note 3.4 we have

$$\{x_a, x_{a+1}, \dots, x_{a+t-1}\} \subset U \quad \text{and} \quad x_{a+t} \notin U. \tag{3.1}$$

Let r be the largest non-negative integer such that $x_{a-i} \in U \mod n$ for $0 \leq i \leq r$ so that

$$\underbrace{x_{a-r-1},}_{\notin U} \underbrace{x_{a-r}, \dots, x_{a-1}, x_a, x_{a+1}, \dots, x_{a+t-1},}_{\in U} \underbrace{x_{a+t}}_{\notin U}. \tag{3.2}$$

It follows that since Γ is an induced subcollection of $\Delta_t(C_n)$ on U, $F_{a-r}, F_{a-r+1}, \dots, F_a \in \Gamma$. We now show that

$$F_i \notin \Gamma \text{ for all } i \notin \{a-r, a-r+1, \dots, a\} \mod n.$$

This follows from the fact that Γ is connected: if any F_i (except for $a-r \leq i \leq a \mod n$) intersects some of the facets $F_{a-r}, \dots, F_a \mod n$, then it must contain x_{a-r-1} or x_{a+t} (as otherwise it would be one of $F_{a-r}, \dots, F_a \mod n$), and hence $F_i \notin \Gamma$.

We have therefore shown that

$$\Gamma = \langle F_{a-r}, F_{a-r+1}, \dots, F_a \rangle \mod n.$$

Next we prove $\Gamma = \Delta_t(L_{|U|})$. Without loss of generality we can assume that $a - r = 1$, so that $\Gamma = \langle F_1, \dots, F_{r+1} \rangle$. Since $\Gamma \neq \Delta_t(C_n)$ we can say that $r + 1 < n$ and therefore we have

$$\text{Vert}(\Gamma) = \{x_1, x_2, \dots, x_{r+t}\}.$$

Since Γ is induced and proper we have $r + t < n$, and therefore we can conclude that

$$\Gamma = \Delta_t(L_{\{x_1,x_2,\ldots,x_{t+r}\}}).$$

□

Example 3.7. Consider the cycle C_7 on vertex set $\mathcal{X} = \{x_1, \ldots x_7\}$ and the simplicial complex $\Delta_4(C_7)$. The following induced subcollections are two runs in $\Delta_4(C_7)$

$$\begin{aligned}\Delta_1 &= \langle\{x_1, x_2, x_3, x_4\}, \{x_2, x_3, x_4, x_5\}\rangle \\ \Delta_2 &= \langle\{x_1, x_2, x_6, x_7\}, \{x_1, x_2, x_3, x_7\}, \{x_1, x_2, x_3, x_4\}\rangle.\end{aligned}$$

Lemma 3.8. *Let* Γ *and* Λ *be two induced subcollections of* $\Delta_t(C_n) = \langle F_1, F_2, \ldots, F_n\rangle$ *which are both composed of runs of lengths* $s_1, \ldots, s_r$*. Then* Γ *and* Λ *are homeomorphic as simplicial complexes. In particular the two simplicial complexes* Γ^c *and* Λ^c *are homeomorphic and have the same reduced homologies.*

Proof. First we suppose $\Delta_t(C_n) = \langle F_1, F_2, \ldots, F_n\rangle$ has standard labeling. If we denote each run of length s_i in Γ and Λ by R_i and R_i', respectively, we have

$$\Gamma = \langle R_1, R_2, \ldots, R_r\rangle \text{ and } \Lambda = \langle {R_1}', {R_2}', \ldots, {R_r}'\rangle$$

where, using the standard labeling, for $1 \leq j_i, h_i \leq n$ we have

$$R_i = \langle F_{j_i}, F_{j_i+1}, \ldots, F_{j_i+s_i-1}\rangle \text{ and } {R_i}' = \langle F_{h_i}, F_{h_i+1}, \ldots, F_{h_i+s_i-1}\rangle \mod n.$$

Then clearly

$$\text{Vert}(\Gamma) = \bigcup_{i=1}^{r} \text{Vert}(R_i) = \bigcup_{i=1}^{r} \{x_{j_i}, x_{j_i+1}, \ldots, x_{j_i+s_i+t-2}\}$$

and

$$\text{Vert}(\Lambda) = \bigcup_{i=1}^{r} \text{Vert}({R_i}') = \bigcup_{i=1}^{r} \{x_{h_i}, x_{h_i+1}, \ldots, x_{h_i+s_i+t-2}\}.$$

Now we define the function $\varphi : \text{Vert}(\Gamma) \longrightarrow \text{Vert}(\Lambda)$ where

$$\varphi(x_{j_i+u}) = x_{h_i+u} \text{ for } 0 \leq u \leq s_i + t - 2.$$

Since φ is a bijective map between vertex set Γ and Λ which preserves faces, we can conclude Γ and Λ are homeomorphic. Therefore, two simplicial complexes Γ^c and Λ^c are homeomorphic as well and have the same reduced homology. □

Therefore, by applying Theorem 2.8 and Lemma 3.8 all the information we need to compute the homologies of induced subcollections of $\Delta_t(C_n)$ depends on the number and the lengths of the runs.

Definition 3.9. For a fixed integer $t \geq 2$, let the pure $(t-1)$-dimensional simplicial complex $\Gamma = \langle F_1, \ldots, F_s\rangle$ be a disjoint union of runs of length $s_1, \ldots, s_r$. Then the sequence of positive integers $s_1, \ldots, s_r$ is called a **run sequence** on $\mathcal{Y} = \text{Vert}(\Gamma)$, and we use the notation

$$E(s_1, \ldots, s_r) = \Gamma^c_{\mathcal{Y}} = \langle (F_1)^c_{\mathcal{Y}}, \ldots, (F_s)^c_{\mathcal{Y}}\rangle.$$

4 Reduced homologies for Betti numbers

Let $I = I_t(C_n)$ be the path ideal of the the cycle C_n for some $t \geq 2$. By applying Hochster's formula (Theorem 2.8), we see that to compute the Betti numbers of R/I, we need to compute the reduced homologies of complements of induced subcollections of Δ. This section is devoted to complex homological calculations. The results here will allow us to compute all Betti numbers of R/I (and more) in the sections that follow. We begin by recalling the Mayer-Vietoris sequence; for more details see for example Hatcher [6] Chapter 2.

Suppose Δ is a simplicial complex and Δ_1 and Δ_2 are two subcollections such that $\Delta = \Delta_1 \cup \Delta_2$. We have the exact sequence of the chain complexes

$$0 \to \mathcal{C}.(\Delta_1 \cap \Delta_2) \to \mathcal{C}.(\Delta_1) \oplus \mathcal{C}.(\Delta_2) \to \mathcal{C}.(\Delta) \to 0$$

which produces the following long exact sequence, called the **Mayer-Vietoris** sequence

$$\cdots \to \widetilde{H}_i(\Delta_1 \cap \Delta_2) \to \widetilde{H}_i(\Delta_1) \oplus \widetilde{H}_i(\Delta_2) \to \widetilde{H}_i(\Delta) \to \widetilde{H}_{i-1}(\Delta_1 \cap \Delta_2) \to \ldots .$$

We make a basic observation.

Lemma 4.1. *Let $E_1, \ldots, E_s$ be subsets of the finite set V where $s \geq 2$ and suppose that $\mathcal{E} = \langle (E_1)^c_V, (E_2)^c_V, ..., (E_s)^c_V \rangle$. Then for any i we have*

i. Suppose $V \setminus \bigcup_{j=2}^{s} E_j \neq \emptyset$. If $\mathcal{E}_1 = \langle (E_1)^c_V \rangle$ and $\mathcal{E}_2 = \langle (E_2)^c_V, ..., (E_s)^c_V \rangle$ then

$$\begin{aligned}\widetilde{H}_i(\mathcal{E}) = \widetilde{H}_i(\mathcal{E}_1 \cup \mathcal{E}_2) &\cong \widetilde{H}_{i-1}(\mathcal{E}_1 \cap \mathcal{E}_2)\\ &= \widetilde{H}_{i-1}(\langle (E_1 \cup E_2)^c_V, ..., (E_1 \cup E_s)^c_V \rangle)\\ &= \widetilde{H}_{i-1}(\langle (E_2)^c_{(V \setminus E_1)}, ..., (E_s)^c_{(V \setminus E_1)} \rangle).\end{aligned}$$

ii. If $E_a \subset E_b$ for some $a \neq b$, then $\mathcal{E} = \langle (E_1)^c_v, \ldots, (\widehat{E_b})^c_V, \ldots, (E_s)^c_V \rangle$.

The decomposition $\mathcal{E} = \mathcal{E}_1 \cup \mathcal{E}_2$ described above is called **standard decomposition** of $\mathcal{E}$.

Proof. The proof of ii is trivial so we shall only prove i.

Since $\mathcal{E}_1$ is a simplex, we have $\widetilde{H}_i(\mathcal{E}_1) = 0$. Also since $V \setminus \bigcup_{i=2}^{s} E_i \neq \emptyset$ we have $\mathcal{E}_2$ is a cone, so from Proposition 2.7 we conclude

$$\widetilde{H}_i(\mathcal{E}_2) = 0 \text{ for all } i.$$

Now $\mathcal{E}_1 \cap \mathcal{E}_2 = \langle (E_1 \cup E_2)^c_V, ..., (E_1 \cup E_s)^c_V \rangle$, so by applying the Mayer-Vietoris sequence we reach the following exact sequence

$$\cdots \longrightarrow \widetilde{H}_i(\mathcal{E}_1) \oplus \widetilde{H}_i(\mathcal{E}_2) \longrightarrow \widetilde{H}_i(\mathcal{E}) \longrightarrow \widetilde{H}_{i-1}(\mathcal{E}_1 \cap \mathcal{E}_2) \longrightarrow \widetilde{H}_{i-1}(\mathcal{E}_1) \oplus \widetilde{H}_{i-1}(\mathcal{E}_2) \longrightarrow \cdots$$

which implies that

$$\begin{aligned}\widetilde{H}_i(\mathcal{E}_1 \cup \mathcal{E}_2) \cong& \widetilde{H}_{i-1}(\mathcal{E}_1 \cap \mathcal{E}_2)\\ =& \widetilde{H}_{i-1}(\langle (E_1 \cup E_2)^c_V, ..., (E_1 \cup E_s)^c_V \rangle).\end{aligned}$$

$\square$

Proposition 4.2. *Let $\Gamma = \langle E_1, \dots, E_s \rangle$ be a pure simplicial complex of dimension $t-1$ over the vertex set $V = \{x_1, \dots, x_n\}$ where $2 \leq t \leq n$. Suppose the connected components of Γ are runs of lengths $s_1, \dots, s_r$, and $\mathcal{E} = E(s_1, \dots, s_r)$. Let $s_j = (t+1)p_j + d_j$ where $p_j \geq 0$ and $0 \leq d_j \leq t$ and $1 \leq j \leq r$. Then for all i, we have*

$i.$ *If $s_j \geq t+2$ then* $\widetilde{H}_i(\mathcal{E})) \cong \widetilde{H}_{i-2}(E(s_1, \dots, s_j - (t+1), \dots, s_r))$;

$ii.$ *If $d_j \neq 1, 2$ then* $\widetilde{H}_i(\mathcal{E}) = 0$*;*

$iii.$ *If $s_j = 2$ and $r \geq 2$ then* $\widetilde{H}_i(\mathcal{E}) = \widetilde{H}_{i-2}(E(s_1, \dots, s_{j-1}, s_{j+1}, \dots, s_r))$*;*

$iv.$ *If $s_j = 1$ and $r \geq 2$ then* $\widetilde{H}_i(\mathcal{E}) = \widetilde{H}_{i-1}(E(s_1, \dots, s_{j-1}, s_{j+1}, \dots, s_r))$*.*

Proof. We assume without loss of generality that $E_1, \dots, E_s$ are ordered such that $E_1, \dots, E_{s_j}$ are the facets of the run of length s_j, and they have standard labeling

$$E_1 = \{x_1, x_2, \dots, x_t\}, E_2 = \{x_2, x_3, \dots, x_{t+1}\}, \dots, E_{s_j} = \{x_{s_j}, x_{s_j+1}, \dots, x_{s_j+t-1}\} \mod n.$$

We have $\mathcal{E} = \langle (E_1)^c_V, (E_2)^c_V, ..., (E_s)^c_V \rangle$. Since $x_1 \in V \backslash \bigcup_{i=2}^{s} E_i$ there is a standard decomposition

$$\mathcal{E} = \langle (E_1)^c_V \rangle \cup \langle (E_2)^c_V, \dots, (E_s)^c_V \rangle.$$

From Lemma 4.1 (i), setting $V' = V \setminus \{x_1, x_2, \dots, x_t\}$, we have

$$\widetilde{H}_i(\mathcal{E}) \cong \widetilde{H}_{i-1}(\langle (E_2)^c_{V'}, \dots, (E_s)^c_{V'} \rangle). \tag{4.1}$$

If $s_j \geq t+2$ from (4.1) we have

$$\widetilde{H}_i(\mathcal{E})) = \widetilde{H}_{i-1}(\langle \{x_{t+1}\}^c_{V'}, (E_{t+2})^c_{V'}, \dots, (E_s)^c_{V'} \rangle) \tag{4.2}$$

and since the following is a standard decomposition

$$\langle \{x_{t+1}\}^c_{V'} \rangle \cup \langle (E_{t+2})^c_{V'}, \dots, (E_{s_j})^c_{V'}, \dots, (E_s)^c_{V'} \rangle$$

from (4.2), Lemma 4.1 (i) and by setting $V'' = V \setminus \{x_1, x_2, \dots, x_{t+1}\}$, we have

$$\widetilde{H}_i(\mathcal{E}) \cong \widetilde{H}_{i-2}(\langle (E_{t+2})^c_{V''}, \dots, (E_{s_j})^c_{V''}, \dots, (E_s)^c_{V''} \rangle).$$

Now the connected components of $\langle E_{t+2}, \dots, E_s \rangle$ are runs of lengths $s_1, \dots, s_j - (t+1), \dots, s_r$, and and therefore we can conclude that for all i

$$\widetilde{H}_i(\mathcal{E}) = \widetilde{H}_{i-2}(E(s_1, \dots, s_j - (t+1), \dots, s_r)).$$

This settles Case (i) of the proposition. Now suppose $1 \leq s_j < t+2$. In this case by (4.1) and Lemma 4.1 (i) and (ii) we see that

$$\widetilde{H}_i(\mathcal{E}) \cong \widetilde{H}_{i-1}(\langle \{x_{t+1}\}^c_{V'}, (E_{s_j+1})^c_{V'}, \dots, (E_s)^c_{V'} \rangle) \quad \text{for all } i. \tag{4.3}$$

1. If $s_j \geq 3$ since $x_{s_j+t-1} \in V' \setminus (\bigcup_{i=s_j+1}^{s} E_i \cup \{x_{t+1}\})$ the simplicial complex

$$\langle \{x_{t+1}\}^c_{V'}, (E_{s_j+1})^c_{V'}, \dots, (E_s)^c_{V'} \rangle$$

is a cone and by Proposition 2.7 and (4.3) we have $\widetilde{H}_i(\mathcal{E}) = 0$ for all i.

2. If $s_j = 2$ and $r \geq 2$, since $x_{t+1} \in V' \setminus (\bigcup_{i=s_j+1}^{s} E_i)$ we have

   $$\langle \{x_{t+1}\}_{V'}^c \rangle \cup \langle (E_{s_j+1})_{V'}^c, \dots, (E_s)_{V'}^c \rangle$$

   is a standard decomposition and then by Lemma 4.1 and (4.3) we have

   $$\begin{aligned} \widetilde{H}_i(\mathcal{E}) &\cong \widetilde{H}_{i-2}(\langle (E_{s_j+1})_{V''}^c, \dots, (E_s)_{V''}^c \rangle) \\ &= \widetilde{H}_{i-2}(E(s_1, \dots, s_{j-1}, s_{j+1}, \dots, s_r)) \quad \text{for all } i. \end{aligned}$$

   This settles Case (iii).

3. If $s_j = 1$ and $r \geq 2$ since $E_1 \cap E_h = \emptyset$ for $1 < h \leq s$, and from (4.1) we have

   $$\widetilde{H}_i(\mathcal{E}) \cong \widetilde{H}_{i-1}(E(s_1, \dots s_{j-1}, s_{j+1}, \dots, s_r)) \text{ for all } i.$$

   This settles Case (iv).

To prove (ii), we use induction on p_j. If $p_j = 0$, then $d_j = s_j \geq 1$. From above we know that $\widetilde{H}_i(\mathcal{E}) = 0$ if $3 \leq s_j \leq t$, and we are done. Now suppose $p_j \geq 1$ and the statement holds for all values less than p_j. We have two cases:

1. If $s_j < t+2$, then since $p_j \geq 1$, we must have $p_j = 1$, $d_j = 0$, and $s_j = t+1$. It was proved above (under the case $3 \leq s_j < t+2$) that $\widetilde{H}_i(\mathcal{E}) = 0$.

2. If $s_j \geq t+2$, by (i) we have

   $$\widetilde{H}_i(\mathcal{E}) \cong \widetilde{H}_{i-2}(E(s_1, \dots, (t+1)(p_j - 1) + d_j, \dots, s_r)) = 0 \text{ when } d_j \neq 1, 2.$$

This proves (ii) and we are done. □

We conclude that for computing the homology of the induced subcollections of path complexes of cycles or lines, the only cases which have to be considered are those of runs of length one or two. We now set about computing these.

Proposition 4.3. *Let* t *and* n *be integers, such that* $2 \leq t \leq n$. *Let* $\alpha, \beta \geq 0$ *and consider*

$$\mathcal{E} = E((t+1)p_1 + 1, \dots, (t+1)p_\alpha + 1, (t+1)q_1 + 2, \dots, (t+1)q_\beta + 2)$$

for nonnegative integers $p_1 \dots, p_\alpha, q_1, \dots, q_\beta$. *Then*

$$\widetilde{H}_i(\mathcal{E}) = \begin{cases} K & i = 2(P+Q) + 2\beta + \alpha - 2 \\ 0 & \textit{otherwise} \end{cases}$$

where $P = \sum_{i=1}^{\alpha} p_i$ *and* $Q = \sum_{i=1}^{\beta} q_i$.

From here on, we use the notation $E(1^\alpha, 2^\beta)$ to denote the complex $\mathcal{E}$ described in the statement of Proposition 4.3 in the case where all the p's and q's are zero; i.e. the case of α runs of length one and β runs of length two.

Proof. First we prove the two cases $\alpha = 0$, $\beta = 1$ and $\alpha = 1$, $\beta = 0$.

1. If $\alpha = 1$, $\beta = 0$, then $\mathcal{E} \cong \langle V \setminus \{x_1, x_2, \dots, x_t\}\rangle = \{\emptyset\}$ where $V = \{x_1, \dots, x_t\}$, and therefore
$$\widetilde{H}_i(\mathcal{E}) = \begin{cases} K & i = -1 \\ 0 & \text{otherwise.} \end{cases}$$
2. If $\alpha = 0$, $\beta = 1$, then $\mathcal{E} \cong \langle (\{x_1, x_2, \dots, x_t\})^c_V, (\{x_2, \dots, x_{t+1}\})^c_V\rangle = \langle \{x_{t+1}\}, \{x_1\}\rangle$ where $V = \{x_1, \dots, x_{t+1}\}$. Since $\mathcal{E}$ is disconnected, we have
$$\widetilde{H}_i(\mathcal{E}) = \begin{cases} K & i = 0 \\ 0 & \text{otherwise.} \end{cases}$$

To prove the statement of the proposition, we use repeated applications of Proposition 4.2 (i), p_1 times to the first run, p_2 times to the second run, and so on till q_β times to the last run as follows.

$$\begin{aligned}
\widetilde{H}_i(\mathcal{E}) &= \widetilde{H}_i(E((t+1)p_1 + 1, \dots, (t+1)p_\alpha + 1, (t+1)q_1 + 2, \dots, (t+1)q_\beta + 2)) \\
&\cong \widetilde{H}_{i-2}(E((t+1)(p_1 - 1) + 1, \dots, (t+1)p_\alpha + 1, (t+1)q_1 + 2, \dots, (t+1)q_\beta + 2)) \\
&\ \ \vdots \\
&\cong \widetilde{H}_{i-2(P+Q)}(E(1^\alpha, 2^\beta)) && \text{apply Proposition 4.2 (iv)} \\
&\cong \widetilde{H}_{i-2(P+Q)-\alpha}(E(2^\beta)) && \text{apply Proposition 4.2 (iii)} \\
&\cong \widetilde{H}_{i-2(P+Q)-\alpha-2\beta+2}(E(2)) && \text{apply Case 2 above} \\
&= \begin{cases} K & i = 2(P+Q) + \alpha + 2\beta - 2 \\ 0 & \text{otherwise.} \end{cases}
\end{aligned}$$

□

An immediate consequence of the above calculations is the homology of the complement of a run, or equivalently, the path complex of any line graph.

Corollary 4.4. *Let t, p and d be integers such that $t \geq 2$, $p \geq 0$, and $0 \leq d \leq t$. Then*
$$\widetilde{H}_i(E((t+1)p + d)) = \begin{cases} K & d = 1,\ i = 2p - 1 \\ K & d = 2,\ i = 2p \\ 0 & \textit{otherwise.} \end{cases}$$

Proof. By Proposition 4.2 (ii), if $d \neq 1, 2$ the homology is zero. In the cases where $d = 1, 2$, the result follows directly from Proposition 4.3. □

We end this section with the calculation of the homology of the complement of the whole path complex of a cycle; this will give us the top degree Betti numbers of the path ideal of a cycle. We will first need a technical lemma.

Lemma 4.5. *Let $R = K[x_1, \dots, x_n]$ be a polynomial ring over a field K, and suppose $\Delta_t(C_n) = \langle F_1, F_2, \dots, F_n\rangle$ is the path complex of a cycle C_n with standard labeling. Let $a, k, s, t \in \{1, \dots, n\}$ be such that $k < t$, and $a + s + t - 1 < n$. Suppose $s = (t+1)p + d$ where $p \geq 0$ and $0 \leq d < t + 1$. Set $V = \{x_a, x_{a+1}, \dots, x_{a+s+t-1}\}$ and*
$$\mathcal{E} = \langle (F_a)^c_V, \dots, (F_{a+s-1})^c_V, \{x_{a+s+t-k}, x_{a+s+t-k+1}, \dots, x_{a+s+t-1}\}^c_V\rangle.$$
Then for all i we have
$$\widetilde{H}_i(\mathcal{E}) = \begin{cases} K & d = 1, i = 2p \\ K & d = k + 1, i = 2p + 1 \\ 0 & \textit{otherwise.} \end{cases}$$

Proof. Without loss of generality we can assume $a = 1$ so that $V = \{x_1, \dots, x_{s+t}\}$ and

$$\mathcal{E} = \langle (F_1)^c_V, \dots, (F_s)^c_V, \{x_{s+t-k+1}, \dots, x_{s+t}\}^c_V \rangle.$$

Since $x_{s+t} \notin F_h$ for $1 \le h \le s$, $\mathcal{E}$ has standard decomposition

$$\mathcal{E} = \langle (F_1)^c_V, (F_2)^c_V, \dots, (F_s)^c_V \rangle \cup \langle (\{x_{s+t-k+1}, x_{s+t-k+2}, \dots, x_{s+t}\})^c_V \rangle$$

and then from Lemma 4.1 (i) and (ii), setting $V_1 = V \setminus \{x_{s+t-k+1}, x_{s+t-k+2}, \dots, x_{s+t}\}$, we have

$$\begin{aligned} \widetilde{H}_i(\mathcal{E}) \cong &\widetilde{H}_{i-1}(\langle (F_1)^c_{V_1}, (F_2)^c_{V_1}, \dots, (F_{s-k})^c_{V_1}, \{x_{s-k+1}, \dots, x_{s+t-k}\}^c_{V_1}, \{x_{s-k+2}, \dots, x_{s+t-k}\}^c_{V_1}, \\ &\quad , \dots, \{x_{s-1}, \dots, x_{s+t-k}\}^c_{V_1}, \{x_s, \dots, x_{s+t-k}\}^c_{V_1} \rangle) \\ = &\widetilde{H}_{i-1}(\langle (F_1)^c_{V_1}, (F_2)^c_{V_1}, \dots, (F_{s-k})^c_{V_1}, \{x_s, \dots, x_{s+t-k}\}^c_{V_1} \rangle) \end{aligned} \tag{4.4}$$

We prove our statement by induction on $|V| = s+t = (t+1)p+d+t$. The base case is $|V| = d+t$, in which case $p = 0$ and $d = s \ge 1$. There are two cases to consider.

1. If $1 \le d \le k$, then $s \le k$, and so by (4.4)

   $$\widetilde{H}_i(\mathcal{E}) = \widetilde{H}_{i-1}(\langle \{x_s, \dots, x_{s+t-k}\}^c_{V_1} \rangle).$$

   The simplex $\{x_s, \dots, x_{s+t-k}\}^c_{V_1}$ is not empty unless $s = d = 1$, and hence we have

   $$\widetilde{H}_i(\mathcal{E}) = \begin{cases} K & d = 1, i = 0 \\ 0 & \text{otherwise.} \end{cases}$$

2. If $d > k$, we use (4.4) to note that since $x_{s-k+t} \notin F_1 \cup \ldots \cup F_{s-k}$, the following is a standard decomposition

   $$\langle (F_1)^c_{V_1}, (F_2)^c_{V_1}, \dots, (F_{s-k})^c_{V_1}, \{x_s, \dots, x_{s+t-k}\}^c_{V_1} \rangle.$$

   Using Lemma 4.1 and (4.4) along with the fact that $s = d \le t$, we find that if $V_2 = V \setminus \{x_s, \dots, x_{s+t}\}$, then

   $$\begin{aligned} \widetilde{H}_i(\mathcal{E}) &\cong \widetilde{H}_{i-2}(\langle \{x_1, \dots, x_{s-1}\}^c_{V_2}, \{x_2, \dots, x_{s-1}\}^c_{V_2}, \dots, \{x_{s-k}, \dots, x_{s-1}\}^c_{V_2} \rangle) \\ &\cong \widetilde{H}_{i-2}(\langle \{x_{s-k}, \dots, x_{s-1}\}^c_{V_2} \rangle). \end{aligned}$$

   Now the simplex $\{x_{s-k}, \dots, x_{s-1}\}^c_{V_2}$ is nonempty unless $s - k = 1$, or in other words, $d = s = k+1$. Therefore

   $$\widetilde{H}_i(\mathcal{E}) = \begin{cases} K & d = k+1, i = 1 \\ 0 & \text{otherwise.} \end{cases}$$

This settles the base case of the induction. Now suppose $|V| = s+t > d+t$ and the theorem holds for all the cases where $|V| < s+t$. Since $|V_1| = (s-k)+t < |V|$ we shall apply (4.4) and use the induction hypothesis on V_1, now with the following parameters: $k_1 = t-k+1$, $s_1 = s-k = (t+1)p+d-k$ and

$$d_1 = \begin{cases} d-k & d \ge k \\ d-k+t+1 & d < k \end{cases} \quad \text{and} \quad p_1 = \begin{cases} p & d \ge k \\ p-1 & d < k. \end{cases}$$

Applying the induction hypothesis on V_1 we see that $\widetilde{H}_i(\mathcal{E}) = 0$ unless one of the following scenarios happen, in which case $\widetilde{H}_i(\mathcal{E}) = K$.

1. $d_1 = 1$ and $i - 1 = 2p_1$.
    (a) When $d \geq k$, this means that $d = k + 1$ and $i = 2p + 1$.
    (b) When $d < k$, this means $d - k + t + 1 = 1$ which implies that $0 \leq d = k - t \leq 0$, and hence $d = 0$ and $t = k$, which is not possible as we have assumed $k < t$.
2. $d_1 = k_1 + 1$ and $i - 1 = 2p_1 + 1$.
    (a) When $d \geq k$, this means that $d - k = t - k + 1 + 1$ and so $d = t + 2$ which is not possible, as we have assumed $d < t + 1$.
    (b) When $d < k$, this means that $d - k + t + 1 = t - k + 1 + 1$ and so $d = 1$ and $i = 2p_1 + 2 = 2p$.

We conclude that $\widetilde{H}_i(\mathcal{E}) = K$ only when $d = 1$ and $i = 2p$, or $d = k + 1$ and $i = 2p + 1$, and $\widetilde{H}_i(\mathcal{E}) = 0$ otherwise. □

Theorem 4.6. *Let $2 \leq t \leq n$ and $\Delta = \Delta_t(C_n)$ be the path complex of a cycle C_n with vertex set $\mathcal{X} = \{x_1, x_2, \dots, x_n\}$. Suppose $n = (t+1)p + d$ where $p \geq 0$, $0 \leq d \leq t$. Then for all i*

$$\widetilde{H}_i(\Delta^c_{\mathcal{X}}) = \begin{cases} K^t & d = 0,\ i = 2p-2,\ p > 0 \\ K & d \neq 0,\ i = 2p-1 \\ 0 & \textit{otherwise.} \end{cases}$$

Proof. If $p = 0$, then $n = t$ and our claim is obvious, so we assume that $p \geq 1$ and therefore $n \geq t+1$. We define the following simplicial complexes

$$\begin{aligned} E_0 &= \langle (F_1)^c_{\mathcal{X}}, (F_2)^c_{\mathcal{X}}, \dots, (F_{n-t+1})^c_{\mathcal{X}} \rangle = E(n-t+1) \\ E_k &= E_{k-1} \cup \langle (F_{n-k+1})^c_{\mathcal{X}} \rangle \text{ for } k = 1, 2, \dots, t-1. \end{aligned} \tag{4.5}$$

Note that $\Delta^c_{\mathcal{X}} = E_{t-1}$. We start with E_0 and apply the Mayer-Vietoris sequence repeatedly to calculate the homologies of the E_k. Since $E_0 = E(n-t+1)$, we find

$$n - t + 1 = (t+1)p + d - t + 1 = \begin{cases} (t+1)p + 1 & d = t \\ (t+1)p & d = t-1 \\ (t+1)(p-1) + d + 2 & d < t-1 \end{cases}$$

which by Corollary 4.4 implies that

$$\widetilde{H}_i(E_0) = \begin{cases} K & d = 0,\ i = 2p-2 \\ K & d = t,\ i = 2p-1 \\ 0 & \text{otherwise.} \end{cases} \tag{4.6}$$

In order to to find the homologies of E_{t-1} we shall recursively apply the Mayer-Vietoris sequence as follows. For a fixed $1 \leq k \leq t-1$ we have the following exact sequence,

$$\widetilde{H}_i(E_{k-1} \cap \langle (F_{n-k+1})^c_{\mathcal{X}} \rangle) \to \widetilde{H}_i(E_{k-1}) \to \widetilde{H}_i(E_k) \to \widetilde{H}_{i-1}(E_{k-1} \cap \langle (F_{n-k+1})^c_{\mathcal{X}} \rangle). \tag{4.7}$$

We claim that for $0 \leq k \leq t-2$,

$$\widetilde{H}_i(E_k \cap \langle (F_{n-k})^c_{\mathcal{X}} \rangle) = \begin{cases} K & d = 0,\ i = 2p-3 \\ K & d = t-k-1,\ i = 2p-2 \\ 0 & \text{otherwise.} \end{cases} \tag{4.8}$$

Setting $\mathcal{X}' = \mathcal{X} \setminus F_{n-k} = \{x_{t-k}, \ldots, x_{n-k-1}\}$ we can write

$$E_k \cap \langle (F_{n-k})^c_{\mathcal{X}} \rangle = \langle (F_1)^c_{\mathcal{X}'}, \ldots, (F_{n-t+1})^c_{\mathcal{X}'}, (F_{n-k+1})^c_{\mathcal{X}'}, \ldots, (F_n)^c_{\mathcal{X}'} \rangle. \tag{4.9}$$

We now compute the $(F_h)^c_{\mathcal{X}'} = \{x_h, \ldots, x_{h+(t-1)}\}^c_{\mathcal{X}'}$ appearing in (4.9).

- When $1 \leq h \leq t-k$ it is clear that
$$(F_h)^c_{\mathcal{X}'} = \{x_{t-k}, x_{t-k+1}, \ldots, x_{t+h-1}\}^c_{\mathcal{X}'}.$$
- When $t-k+1 \leq h \leq n-t-k-1$ then $2t-k \leq h+t-1 \leq n-k-2$, and so
$$(F_h)^c_{\mathcal{X}'} = \{x_{t-k}, \ldots, x_{h-1}, x_{h+t}, \ldots, x_{n-k-1}\}.$$
- When $n-k-t \leq h \leq n-t+1$, then $n-k-1 \leq h+t-1 \leq n$, and therefore
$$(F_h)^c_{\mathcal{X}'} = \{x_h, \ldots, x_{n-k-1}\}^c_{\mathcal{X}'}.$$
- When $0 \leq j \leq k-1$, then $t-k \leq -j+(t-1) \leq t-1$ and so we have
$$F_{n-j} = \{x_{n-j}, \ldots, x_{n-j+(t-1)}\} = \{x_{n-j}, \ldots, x_n, x_1, \ldots, x_{t-j-1}\} \mod n$$
which implies that
$$(F_{n-j})^c_{\mathcal{X}'} = \{x_{t-k}, x_{t-k+1}, \ldots, x_{t-j-1}\}^c_{\mathcal{X}'}.$$

From the observations above, (4.9) and Lemma 4.1 (ii) we see that

$$E_k \cap \langle (F_{n-k})^c_{\mathcal{X}} \rangle = \langle \{x_{t-k}\}, F_{t-k+1}, \ldots, F_{n-t-k-1}, \{x_{n-t+1}, \ldots, x_{n-k-1}\} \rangle^c_{\mathcal{X}'}. \tag{4.10}$$

We now consider the following scenarios.

1. Suppose $p = 1$. In this situation, $n = t+d+1 \leq 2t+1$ which implies that $n-t-k-1 \leq t-k$. Therefore, (4.10) becomes
$$E_k \cap \langle (F_{n-k})^c_{\mathcal{X}} \rangle = \langle \{x_{t-k}\}, \{x_{n-t+1}, \ldots, x_{n-k-1}\} \rangle^c_{\mathcal{X}'}. \tag{4.11}$$
   (a) If $d \leq t-k-2$, then $n-t+1 = t+d+1-t+1 = d+2 \leq t-k$. As well, since $n \geq t+1$, we have $n-k-1 \geq t-k$. It follows that in this situation, $x_{t-k} \in \{x_{n-t+1}, \ldots, x_{n-k-1}\}$ which means that (4.11) becomes $E_k \cap \langle (F_{n-k})^c_{\mathcal{X}} \rangle = \langle \{x_{t-k}\}^c_{\mathcal{X}'} \rangle$. Also note that $\mathcal{X}' = \{x_{t-k}\}$ only when $d = 0$. It follows that
$$\widetilde{H}_i(E_k \cap \langle (F_{n-k})^c \rangle) \cong \begin{cases} K & d = 0,\ i = -1 \\ 0 & \text{otherwise.} \end{cases}$$
   (b) If $d > t-k-2$. In this situation, $x_{t-k} \notin \{x_{n-t+1}, \ldots, x_{n-k-1}\}$ which means that we can apply Lemma 4.1 (i), with $\mathcal{X}'' = \mathcal{X}' \setminus \{x_{t-k}\}$ to find that for all i
$$\widetilde{H}_i(E_k \cap \langle (F_{n-k})^c \rangle) = \widetilde{H}_{i-1}(\langle \{x_{n-t+1}, \ldots, x_{n-k-1}\} \rangle^c_{\mathcal{X}''}).$$
   Moreover, $\{x_{n-t+1}, \ldots, x_{n-k-1}\} = \mathcal{X}''$ only when $d = t-k-1$, and so we have
$$\widetilde{H}_i(E_k \cap \langle (F_{n-k})^c \rangle) \cong \begin{cases} K & d = t-k-1,\ i = 0 \\ 0 & \text{otherwise.} \end{cases}$$

2. Suppose $p \geq 2$. In this case it is easy to see that $n-t-k-1 > t-k$ and $n-t+1 > t-k$. Therefore, we can apply Lemma 4.1 (i) with $\mathcal{X}'' = \mathcal{X}' \setminus \{x_{t-k}\}$ to (4.10) to conclude that for all i

   $$\widetilde{H}_i(E_k \cap \langle (F_{n-k})^c_{\mathcal{X}} \rangle) = \widetilde{H}_{i-1}(\langle F_{t-k+1}, \dots, F_{n-t-k-1}, \{x_{n-t+1}, \dots, x_{n-k-1}\} \rangle^c_{\mathcal{X}''}).$$

   Now we use Lemma 4.5 with values $a = t-k+1$ and $s = n-2t-1 = (p-2)(t+1)+d+1$ to conclude that

   $$\widetilde{H}_i(E_k \cap \langle (F_{n-k})^c_{\mathcal{X}} \rangle) = \begin{cases} K & d = 0,\ i = 2p-3 \\ K & d = t-k-1,\ i = 2p-2 \\ 0 & \text{otherwise} \end{cases}$$

This settles (4.8). We now return to finding $\widetilde{H}_i(E_{t-1})$ by recursively using the Mayer-Vietoris sequence to find $\widetilde{H}_i(E_k)$.

1. If $0 < d < t$ then by (4.8) we know that $\widetilde{H}_i(E_{k-1} \cap \langle (F_{n-k+1})^c_{\mathcal{X}} \rangle)$ is nonzero only when $i = 2p-2$ and $k = t-d$. We apply this observation and (4.6) to the exact sequence (4.7) to see that

   $$\widetilde{H}_i(E_k) = \widetilde{H}_i(E_{k-1}) = \widetilde{H}_i(E_0) = 0 \text{ for } 1 \leq k \leq t-d-1.$$

   Once again we use (4.7) to observe that

   $$\widetilde{H}_i(E_k) = \begin{cases} 0 & 1 \leq k \leq t-d-1 \\ \widetilde{H}_{i-1}(E_{k-1} \cap \langle (F_{n-k+1})^c \rangle) & k = t-d \\ \widetilde{H}_i(E_{t-d}) & t-d < k \leq t-1. \end{cases}$$
   $$= \begin{cases} K & k \geq t-d,\ i = 2p-1 \\ 0 & \text{otherwise.} \end{cases}$$

   We can conclude that in this case

   $$\widetilde{H}_i(E_{t-1}) = \begin{cases} K & i = 2p-1 \\ 0 & \text{otherwise.} \end{cases}$$

2. If $d = t$ then by (4.8) we know that $\widetilde{H}_i(E_{k-1} \cap \langle (F_{n-k+1})^c_{\mathcal{X}} \rangle)$ is always zero. We apply this fact along with (4.6) to the sequence in (4.7) to observe that

   $$\widetilde{H}_i(E_k) \cong \widetilde{H}_i(E_0) = \begin{cases} K & i = 2p-1 \\ 0 & \text{otherwise} \end{cases} \quad \text{for } k \in \{1, 2, \dots, t-1\}.$$

3. If $d = 0$ then by (4.8) we know that $\widetilde{H}_i(E_{k-1} \cap \langle (F_{n-k+1})^c_{\mathcal{X}} \rangle)$ is zero unless $i = 2p-3$, and by (4.6) we know $\widetilde{H}_i(E_0)$ is zero unless $i = 2p-2$. Applying these facts to (4.7) we see that

   $$\widetilde{H}_i(E_k) = \widetilde{H}_i(E_0) = 0 \quad \text{for } i \neq 2p-2.$$

   When $i = 2p-2$, the sequence (4.7) produces an exact sequence

   $$0 \longrightarrow \overbrace{\widetilde{H}_{2p-2}(E_0)}^{K} \longrightarrow \widetilde{H}_{2p-2}(E_1) \longrightarrow \overbrace{\widetilde{H}_{2p-3}(E_0 \cap \langle (F_n)^c \rangle)}^{K} \longrightarrow 0.$$

Therefore

$$\widetilde{H}_i(E_1) = \begin{cases} K^2 & i = 2p-2 \\ 0 & \text{otherwise.} \end{cases}$$

We repeat the above method, recursively , for values $k = 2, 3, \ldots, t-1$

$$0 \longrightarrow \overbrace{\widetilde{H}_{2p-2}(E_{k-1})}^{K^k} \longrightarrow \widetilde{H}_{2p-2}(E_k) \longrightarrow \overbrace{\widetilde{H}_{2p-3}(E_{k-1} \cap \langle (F_{n-k+1})^c \rangle)}^{K} \longrightarrow 0$$

and conclude that for $1 \leq k \leq t-1$

$$\widetilde{H}_i(E_k) = \begin{cases} K^{k+1} & i = 2p-2 \\ 0 & \text{otherwise.} \end{cases}$$

We put this all together

$$\widetilde{H}_i(E_{t-1}) = \begin{cases} K^t & d = 0,\ i = 2p-2,\ p > 0 \\ K & d \neq 0,\ i = 2p-1 \\ 0 & \text{otherwise} \end{cases}$$

and this proves the statement of the theorem. $\square$

5 Betti numbers of path ideals

We are now ready to apply the homological calculations from the previous section, and compute Betti numbers of path ideals. If I is the degree t path ideal of a cycle or line, then

$$\beta_{i,j}(R/I) = 0 \text{ for all } i \geq 1 \text{ and } j > ti; \tag{5.1}$$

see for example [8] 3.3.4. By Theorem 2.8, to compute the Betti numbers of I of degree less than n, we should consider the complements of proper induced subcollection of $\Delta = \Delta_t(C_n)$ or $\Delta_t(L_n)$ and, for degree n we should consider Δ^c. We first compute the $\mathbb{N}$-graded Betti numbers of degree n of path ideals.

Theorem 5.1 (**Betti numbers of degree** n)**.** *Let* p*,* t*,* n*,* d *be integers such that* $n = (t+1)p + d$*, where* $p \geq 0$*,* $0 \leq d \leq t$*, and* $2 \leq t \leq n$*. If* C_n *is a cycle over* n *vertices, then*

$$\beta_{i,n}(R/I_t(C_n)) = \begin{cases} t & d = 0,\ i = 2\left(\dfrac{n}{t+1}\right) \\[2ex] 1 & d \neq 0,\ i = 2\left(\dfrac{n-d}{t+1}\right) + 1 \\[2ex] 0 & \textit{otherwise.} \end{cases}$$

Proof. Suppose $\Delta = \Delta_t(C_n)$. By Theorem 2.8 $\beta_{i,n}(R/I_t(C_n)) = \dim_K \widetilde{H}_{i-2}(\Delta^c_{\mathcal{X}})$ and the result now follows directly from Theorem 4.6. $\square$

We now focus on Betti numbers of degree less than n. From Hochster's formula we can conclude that this comes down to counting induced subcollections of certain kinds. The next theorem, which is essential to our further discussions, is similar to statement proved for the edge ideal of a cycle in [8].

Definition 5.2. Let i and j be positive integers. We call an induced subcollection Γ of $\Delta_t(C_n)$ an (i,j)**-eligible subcollection** of $\Delta_t(C_n)$ if Γ is composed of disjoint runs of lengths

$$(t+1)p_1+1,\dots,(t+1)p_\alpha+1,(t+1)q_1+2,\dots,(t+1)q_\beta+2 \tag{5.2}$$

for nonnegative integers $\alpha,\beta,p_1,p_2,\dots,p_\alpha,q_1,q_2,\dots,q_\beta$, which satisfy the following conditions

$$\begin{array}{rcl} j &=& (t+1)(P+Q)+t(\alpha+\beta)+\beta \\ i &=& 2(P+Q)+2\beta+\alpha, \end{array}$$

where $P=\sum_{i=1}^{\alpha} p_i$ and $Q=\sum_{i=1}^{\beta} q_i$.

The reason for this terminology will become clear in the following statement.

Theorem 5.3. *Let $I=I(\Lambda)$ be the facet ideal of an induced subcollection Λ of $\Delta_t(C_n)$. Suppose i and j are integers with $i\le j<n$. Then the $\mathbb{N}$-graded Betti number $\beta_{i,j}(R/I)$ is the number of (i,j)-eligible subcollections of Λ.*

Proof. Since $\Delta(I)=\Lambda$ from Theorem 2.8 we have

$$\beta_{i,j}(R/I)=\sum_{\Gamma\subset\Lambda,|\mathrm{Vert}(\Gamma)|=j}\dim_K \widetilde{H}_{i-2}(\Gamma^c_{\mathrm{Vert}(\Gamma)})$$

where Vert(Γ) is the vertex set of Γ and the sum is taken over induced subcollections Γ of Λ.

Each induced subcollection of Λ is clearly an induced subcollection of $\Delta_t(C_n)$, and can therefore be written as a disjoint union of runs. So from Proposition 4.2 we can conclude the only Γ whose complements have nonzero homology are those corresponding to run sequences of the form (5.2). Such subcollections have j vertices where by Definition 3.5

$$\begin{array}{rcl} j &=& ((t+1)p_1+t)+\cdots+((t+1)p_\alpha+t)+((t+1)q_1+t+1)\cdots+((t+1)q_\beta+t+1) \\ &=& (t+1)(P+Q)+t(\alpha+\beta)+\beta. \end{array} \tag{5.3}$$

So

$$\Gamma^c_{\mathrm{Vert}(\Gamma)}=E((t+1)p_1+1,\dots,(t+1)p_\alpha+1,(t+1)q_1+2,\dots,(t+1)q_\beta+2)$$

and by Proposition 4.3 we have

$$\dim_K(\widetilde{H}_{i-2}(\Gamma^c_{\mathrm{Vert}(\Gamma)}))=\begin{cases} 1 & i=2(P+Q)+2\beta+\alpha \\ 0 & \text{otherwise.} \end{cases} \tag{5.4}$$

From (5.3) and (5.4) we see that each induced subcollection Γ corresponding to a run sequence as in (5.2) contributes 1 unit to $\beta_{i,j}$ if and only if

$$\begin{array}{rcl} j &=& (t+1)(P+Q)+t(\alpha+\beta)+\beta \\ i &=& 2(P+Q)+2\beta+\alpha. \end{array}$$

□

The following corollary is a special case of Theorem 5.3.

Corollary 5.4. *Let $I=I(\Lambda)$ be the facet ideal of an induced subcollection Λ of $\Delta_t(C_n)$. Then for every i, $\beta_{i,ti}(R/I)$, is the number of induced subcollections of Λ which are composed of i runs of length 1.*

Proof. From Theorem 5.3 we have $\beta_{i,ti}(R/I)$ is the number of (i, ti)-eligible subcollections of Λ. With notation as in Definition 5.2 we have

$$\begin{cases} ti = (t+1)(P+Q) + t(\alpha+\beta) + \beta \\ i = 2(P+Q) + (\alpha+\beta) + \beta \end{cases} \Rightarrow ti = 2t(P+Q) + t(\alpha+\beta) + t\beta$$

Putting the two equations for ti together, we conclude that $(t-1)(P+Q+\beta) = 0$. But β, P, $Q \geq 0$ and $t \geq 2$, so we must have

$$\beta = P = Q = 0 \Rightarrow p_1 = p_2 = \cdots = p_\alpha = 0.$$

So $\alpha = i$ and Γ is composed of i runs of length one. □

Theorem 5.3 holds in particular for $\Lambda = \Delta_t(L_m)$ and $\Lambda = \Delta_t(C_n)$ for any integers m, n. Our next statement is in a sense a converse to Theorem 5.3.

Proposition 5.5. *Let t and n be integers such that $2 \leq t \leq n$ and $I = I(\Lambda)$ be the facet ideal of Λ where Λ is an induced subcollection of $\Delta_t(C_n)$. Then for each $i, j \in \mathbb{N}$ with $i \leq d < n$, if $\beta_{i,j}(R/I) \neq 0$, there exist nonnegative integers ℓ, d such that*

$$\begin{cases} i &= \ell + d \\ j &= t\ell + d \end{cases}$$

Proof. From Theorem 5.3 we know $\beta_{i,j}$ is equal to the number of (i, j)-eligible subcollections of Λ, where with notation as in Definition 5.2 we have

$$\begin{cases} j = (t+1)(P+Q) + t(\alpha+\beta) + \beta \\ i = 2(P+Q) + (\alpha+\beta) + \beta. \end{cases}$$

It follows that

$$j - i = (t-1)(P+Q+\alpha+\beta) \quad \text{and} \quad ti - j = (t-1)(P+Q+\beta). \tag{5.5}$$

We now show that there exist positive integers ℓ, d such that $i = \ell + d$ and $j = t\ell + d$.

$$\begin{cases} i = \ell + d \\ j = t\ell + d \end{cases} \Rightarrow \ \ell = \frac{j-i}{t-1} \quad \text{and} \quad d = \frac{ti-j}{t-1} \ .$$

From (5.5) we can see that i and j as described above are nonnegative integers. □

Theorem 5.3 tells us that to compute Betti numbers of induced subcollections of $\Delta_t(C_n)$ we need to count the number of its induced subcollections which consist of disjoint runs of lengths one and two. The next few pages are dedicated to counting such subcollections. We use some combinatorial methods to generalize a helpful formula which can be found in Stanley's book [10] on page 73.

Lemma 5.6. *Consider a collection of n points arranged on a line. The number of ways of coloring k points, when there are at least t uncolored points on the line between each colored point is*

$$\binom{n-(k-1)t}{k}.$$

Proof. First label the points from $1, 2, \dots, n$ from left to right, and let $a_1 < a_2 < \cdots < a_k$ be the colored points. For $1 \leq i \leq k-1$, we define x_i to be the number of points, including a_i, which are between a_i and a_{i+1}, and x_0 to be the number of points which exist before a_1, and x_k the number of points, including a_k, which are after a_k.



If we consider the sequence $x_0, x_1, \dots, x_k$ it is not difficult to see that there is a one to one correspondence between the positive integer solutions of the following equation and the ways of coloring k points of n points on a line with at least t uncolored points between each two colored points.

$$x_0 + x_1 + \cdots + x_k = n \quad x_0 \geq 0, x_i > t, \text{ for } 1 \leq i \leq k-1, \text{ and } x_k \geq 1.$$

So we only need to find the number of positive integer solutions of this equation. Consider the following equation

$$(x_0 + 1) + (x_1 - t) + \cdots + (x_{k-1} - t) + x_k = n - (k-1)t + 1$$

where $x_0 + 1 \geq 1$, $x_i - t \geq 1$, for $i = 0 \dots, k-1$ and $x_k \geq 1$. The number of positive integer solution of this equation is (see for example [5] page 29)

$$\binom{n-(k-1)t}{k}.$$

□

Corollary 5.7. *Let C_n be a graph cycle and with the standard labeling let Γ be a proper subcollection of $\Delta_t(C_n)$ with k facets $F_a, \dots, F_{a+k-1} \mod n$. The number of induced subcollections of Γ which are composed of m runs of length one is*

$$\binom{k-(m-1)t}{m}.$$

Proof. To compute the number of induced subcollections of Γ which are composed of m runs of length one, it is enough to consider the facets $F_a, \dots, F_{a+k-1}$ as points arranged on a line and compute the number of ways which we can color m points of these k arranged points with at least t uncolored points between each two consecutive colored points. Therefore, by Lemma 5.6 we have the number of induced subcollections of Γ which are composed of m runs of length one is $\binom{k-(m-1)t}{m}$. □

Proposition 5.8. *Let C_n be a graph cycle with vertex set $\mathcal{X} = \{x_1, \dots, x_n\}$. The number of induced subcollections of $\Delta_t(C_n)$ which are composed of m runs of length one is*

$$\frac{n}{n-mt}\binom{n-mt}{m}.$$

Proof. Recall that $\Delta_t(C_n) = \langle F_1, \dots, F_n \rangle$ with standard labeling. First we compute the number of induced subcollections of $\Delta_t(C_n)$ which consist of m runs of length one and do not contain the vertex x_n. There are t facets of $\Delta_t(C_n)$ which contain x_n, the remaining facets are $F_1, \dots, F_{n-t}$, and so by Corollary 5.7 the number we are looking for is

$$\binom{n-t-(m-1)t}{m} = \binom{n-mt}{m}. \tag{5.6}$$

Now we are going to compute the number of induced subcollections Γ which consist of m runs of length one and include x_n. We have t facets which contain x_n, they are $F_{n-t+1}\ldots,F_n$. Each such Γ will contain one $F_i \in \{F_{n-t+1}\ldots,F_n\}$ as the run containing x_n, and $m-1$ other runs of length one which have to be chosen so that they are disjoint from F_i. So we are looking for $m-1$ runs of length one in the subcollection $\Gamma' = \langle F_{i+t},\ldots,F_{i-t}\rangle \mod n$. The subcollection Γ' has $n-2t-1$ facets, so by Corollary 5.7 it has

$$\binom{n-2t-1-(m-2)t}{m-1} = \binom{n-mt-1}{m-1}$$

induced subcollections that consist of runs of length one. Putting this together with the number of ways to choose F_i and with (5.6) we conclude that the number of induced subcollections of $\Delta_t(C_n)$ which are composed of m runs of length one is

$$t\binom{n-mt-1}{m-1} + \binom{n-mt}{m} = \frac{n}{n-mt}\binom{n-mt}{m}.$$

□

We apply these counting facts to find Betti numbers in specific degrees; the formula in (iii) below (that of a line graph) was also computed by Bouchat, Ha and O'Keefe [1] using Eliahou-Kervaire techniques.

Corollary 5.9. *Let $n \geq 2$ and t be an integer such that $2 \leq t \leq n$. Then we have*

i. For the cycle C_n we have

$$\beta_{i,it}(R/I_t(C_n)) = \frac{n}{n-it}\binom{n-it}{i}.$$

ii. For any proper induced subcollection Λ of $\Delta_t(C_n)$ with k facets we have

$$\beta_{i,ti}(R/I(\Lambda) = \binom{k-(i-1)t}{i}.$$

iii. For the line graph L_n, we have

$$\beta_{i,ti}(R/I_t(L_n) = \binom{n-it+1}{i}.$$

Proof. From Corollary 5.4 we have $\beta_{i,it}(R/I)$ in each of the three cases (i), (ii) and (iii) is the number of induced subcollections of $\Delta_t(C_n)$, Λ and $\Delta_t(L_n)$, respectively, which are composed of i runs of length 1. Case (i) now follows from Proposition 5.8, while (ii) and (iii) follow directly from Corollary 5.7. □

The following Lemma is the core of our counting later on in this section.

Lemma 5.10. *Let $\Delta_t(C_n) = \langle F_1,F_2,\ldots,F_n\rangle$, $2 \leq t \leq n$, be the standard labeling of the path complex of a cycle C_n on vertex set $\mathcal{X} = \{x_1,\ldots,x_n\}$. Let i be a positive integer and $\Gamma = \langle F_{c_1},F_{c_2},\ldots,F_{c_i}\rangle$ be an induced subcollection of $\Delta_t(C_n)$ consisting of i runs of length 1, with $1 \leq c_1 < c_2 < \cdots < c_i \leq n$. Suppose Σ is the induced subcollection on Vert$(\Gamma) \cup \{x_{c_u+t}\}$ for some $1 \leq u \leq i$. Then*

$$|\Sigma| = \begin{cases} |\Gamma|+t & u<i \text{ and } c_{u+1} = c_u+t+1 \\ |\Gamma|+1 & u=i \text{ or } c_{u+1} > c_u+t+1 \end{cases}$$

Proof. Since Γ consists of runs of length one and each $F_{c_u} = \{x_{c_u}, x_{c_u+1}, \ldots, x_{c_u+t-1}\}$ we must have $c_{u+1} > c_u + t \mod n$ for $u \in \{1, 2, \ldots, i-1\}$. There are two ways that x_{c_u+t} could add facets to Γ to obtain Σ.

1. If $c_{u+1} = c_u + t + 1$ then $F_{c_u}, F_{c_u+1}, \ldots, F_{c_u+t+1} = F_{c_{u+1}} \in \Sigma$ or in other words, we have added t new facets to Γ.
2. If $c_{u+1} > c_u + t + 1$ or $u = i$ then $F_{c_u+1} \in \Sigma$, and therefore one new facet is added to Γ.

□

The following propositions, which generalize Lemma 7.4.22 in [8], will help us compute the remaining Betti numbers.

Proposition 5.11. *Let $\Delta_t(C_n) = \langle F_1, F_2, \ldots, F_n \rangle$, $2 \leq t \leq n$, be the standard labeling of the path complex of a cycle C_n on vertex set $\mathcal{X} = \{x_1, \ldots, x_n\}$. Also let i, j be positive integers such that $j \leq i$ and $\Gamma = \langle F_{c_1}, F_{c_2}, \ldots, F_{c_i} \rangle$ be an induced subcollection of $\Delta_t(C_n)$ consisting of i runs of length 1, with $1 \leq c_1 < c_2 < \cdots < c_i \leq n$. Suppose $W = Vert(\Gamma) \cup A \subsetneq \mathcal{X}$ for some subset A of $\{x_{c_1+t}, \ldots, x_{c_i+t}\} \mod n$ with $|A| = j$. Then the induced subcollection Σ of $\Delta_t(C_n)$ on W is an $(i+j, ti+j)$-eligible subcollection.*

Proof. Since Γ consists of runs of length one and each $F_{c_u} = \{x_{c_u}, x_{c_u+1}, \ldots, x_{c_u+t-1}\}$ we must have $c_{u+1} > c_u + t \mod n$ for $u \in \{1, 2, \ldots, i-1\}$. The runs (or connected components) of Σ are of the form $\Sigma' = \Sigma_U$ where $U \subseteq W$, and can have one of the following possible forms.

a. For some $a \leq i$:
$$U = F_{c_a},$$
and therefore $\Sigma' = \langle F_{c_a} \rangle$ is a run of length 1.

b. For some $a \leq i$:
$$U = F_{c_a} \cup \{x_{c_a+t}\},$$
and therefore $c_{a+1} > c_a + t + 1$, so from Lemma 5.10 we have $\Sigma' = \langle F_{c_a}, F_{c_a+1} \rangle$ is a run of length 2.

c. For some $a \leq i$:
$$U = F_{c_a} \cup F_{c_{a+1}} \cup \cdots \cup F_{c_{a+r}} \cup \{x_{c_a+t}, x_{c_{a+1}+t}, \ldots, x_{c_{a+r-1}+t}\} \mod n$$
and $F_{c_{a+j}} = F_{c_a+j(t+1)}$ for $j = 0, 1, \ldots, r$ and $r \geq 1$. Then from Lemma 5.10 above we know Σ' is a run of length $r + 1 + tr = (t+1)r + 1$.

d. For some $a \leq i$:
$$U = F_{c_a} \cup F_{c_{a+1}} \cup \cdots \cup F_{c_{a+r}} \cup \{x_{c_a+t}, x_{c_{a+1}+t}, \ldots, x_{c_{a+r}+t}\} \mod n$$
and $F_{c_{a+j}} = F_{c_a+j(t+1)}$ for $j = 0, 1, \ldots, r$ and $r \geq 1$, and $c_{a+r+1} > c_{a+r}+t+1$ or $a + r = i$. Then from Lemma 5.10 we have Σ' is a run of length $r + 1 + tr + 1 = (t+1)r + 2$.

So we have shown that Σ consists of runs of length 1 and 2 $\mod t+1$.

Suppose the runs in Σ are of the form described in (5.2). By Definition 3.5 we have

$$\begin{aligned}|\text{Vert}(\Sigma)| &= (t+1)p_1 + t + \cdots + (t+1)p_\alpha + t + (t+1)q_1 + t + 1 + \cdots + (t+1)q_\beta + t + 1\\ &= (t+1)P + t\alpha + (t+1)Q + t\beta + \beta\\ &= (t+1)(P+Q) + t(\alpha+\beta) + \beta.\end{aligned}$$

On the other hand by the definition of Σ we know that, Σ has $ti+j$ vertices and therefore

$$ti + j = (t+1)(P+Q) + t(\alpha+\beta) + \beta.$$

It remains to show that $i+j = 2(P+Q) + (\alpha+\beta) + \beta$. Note that if $j=0$ then $\beta = P = Q = 0$ and hence

$$j = 0 \implies P + Q + \beta = 0. \tag{5.7}$$

Moreover each vertex $x_{c_v+t} \in A$ either increases the length of a run in Γ by one and hence increases β (the number of runs of length 2 in Γ) by one, or increases the length of a run by $t+1$, in which case $P+Q$ increases by 1. We can conclude that if we add j vertices to Γ, $P+Q+\beta$ increases by j. From this and (5.7) we have $j = P+Q+\beta$. Now we solve the following system

$$\left\{\begin{array}{rcl} ti+j &=& (t+1)(P+Q)+t(\alpha+\beta)+\beta \\ j &=& P+Q+\beta \end{array}\right. \begin{array}{l} \implies ti = t(P+Q)+t(\alpha+\beta) \\ \implies i = P+Q+\alpha+\beta \end{array}$$

$$\implies \left\{\begin{array}{rcl} i &=& P+Q+\alpha+\beta \\ j &=& P+Q+\beta \end{array}\right. \implies i+j = 2(P+Q)+(\alpha+\beta)+\beta.$$

□

Proposition 5.12. *Let C_n be a cycle, $2 \le t \le n$, and i and j be positive integers. Suppose Σ is an $(i+j, ti+j)$-eligible subcollection of $\Delta_t(C_n)$, $2 \le t \le n$. Then with notation as in Definition 5.2, there exists a unique induced subcollection Γ of $\Delta_t(C_n)$ of the form $\langle F_{c_1}, F_{c_2}, \ldots, F_{c_i}\rangle$ with $1 \le c_1 < c_2 < \cdots < c_i \le n$ consisting of i runs of length 1, and a subset A of $\{x_{c_1+t}, \ldots, x_{c_i+t}\} \mod n$, with $|A| = j$ such that $\Sigma = \Delta_t(C_n)_W$ where $W = \text{Vert}(\Gamma) \cup A$.*

Moreover if $\mathcal{R} = \langle F_h, F_{h+1}, \ldots, F_{h+m}\rangle \mod n$ is a run in Σ with $|\mathcal{R}| = 2 \mod (t+1)$, then $F_{h+m} \notin \Gamma \mod n$.

Proof. Suppose Σ consists of runs $R'_1, R'_2, \ldots, R'_{\alpha+\beta}$ where for $k = 1, 2, \ldots, \alpha+\beta$

$$\begin{aligned} R'_k &= \langle F_{h_k}, F_{h_k+1}, \ldots, F_{h_k+m_k-1}\rangle && \mod n\\ \text{Vert}(R'_k) &= \{x_{h_k}, x_{h_k+1}, \ldots, x_{h_k+m_k+t-2}\} && \mod n\\ h_{k+1} &\ge t + h_k + m_k && \mod n\end{aligned}$$

and

$$m_k = \left\{\begin{array}{ll} (t+1)p_k + 1 & \text{for} \quad k = 1, 2, \ldots, \alpha \\ (t+1)q_{k-\alpha} + 2 & \text{for} \quad k = \alpha+1, \alpha+2, \ldots, \alpha+\beta. \end{array}\right. \tag{5.8}$$

For each k, we remove the following vertices from Vert(R'_k)

$$\begin{array}{lll} x_{h_k+t}, x_{h_k+2t+1}, \dots, x_{h_k+p_k t+(p_k-1)} & \mod n & \text{if } 1 \leq k \leq \alpha \text{ and } p_k \neq 0 \\ x_{h_k+t}, x_{h_k+2t+1}, \dots, x_{h_k+(q_{k-\alpha}+1)t+q_{k-\alpha}} & \mod n & \text{if } \alpha+1 \leq k \leq \alpha+\beta \end{array} \tag{5.9}$$

Let $\Gamma = \langle R_1, R_2, \dots R_{\alpha+\beta} \rangle$ be the induced subcollection on the remaining vertices of Σ, where

$$R_k = \begin{cases} \langle F_{h_k}, F_{h_k+t+1}, \dots, F_{h_k+(t+1)p_k} \rangle & \mod n \quad \text{for } 1 \leq k \leq \alpha \\ \langle F_{h_k}, F_{h_k+t+1}, \dots, F_{h_k+(t+1)q_{k-\alpha}} \rangle & \mod n \quad \text{for } \alpha+1 \leq k \leq \alpha+\beta. \end{cases} \tag{5.10}$$

In other words, $\mod n$, Γ has facets

$$F_{h_1}, F_{h_1+t+1}, \dots, F_{h_1+(t+1)p_1}, F_{h_2}, F_{h_2+t+1}, \dots, F_{h_2+(t+1)p_2}, \dots, F_{h_{\alpha+\beta}}, \dots, F_{h_{\alpha+\beta}+(t+1)q_\beta}.$$

It is clear that each R_k consists of runs of length one. Since Γ is a subcollection of Σ, no runs of R_k and $R_{k'}$ are connected to one another if $k \neq k'$, and hence we can conclude Γ is an induced subcollection of $\Delta_t(C_n)$ which is composed of runs of length one. From (5.10) we have the number of runs of length 1 in Γ (or the number of facets of Γ) is equal to

$$(p_1+1)+(p_2+1)+\dots+(p_\alpha+1)+(q_1+1)+\dots+(q_\beta+1) = P+Q+\alpha+\beta = i.$$

Therefore, Γ is an induced subcollection of $\Delta_t(C_n)$ which is composed of i runs of length 1. We relabel the facets of Γ as $\Gamma = \langle F_{c_1}, \dots, F_{c_i} \rangle$. Now consider the following subset of $\{x_{c_1+t}, \dots, x_{c_i+t}\}$ as A

$$\bigcup_{k=1, p_k \neq 0}^{\alpha} \{x_{h_k+t}, x_{h_k+2t+1}, \dots, x_{h_k+p_k t+(p_k-1)}\} \cup \bigcup_{k=\alpha+1}^{\alpha+\beta} \{x_{h_k+t}, x_{h_k+2t+1}, \dots, x_{h_k+(q_{k-\alpha}+1)t+q_{k-\alpha}}\}$$

by (5.9) we have:

$$|A| = (p_1+p_2+\dots+p_\alpha)+(q_1+1\dots+q_\beta+1) = P+Q+\beta = j.$$

Then if we set

$$W = (\bigcup_{h=1}^{i} F_{c_h}) \cup A$$

we clearly have $\Sigma = (\Delta_t(C_n))_W$. This proves the existence of Γ, we now prove its uniqueness. Let $\Lambda = \langle F_{s_1}, F_{s_2}, \dots, F_{s_i} \rangle$ be an induced subcollection of $\Delta_t(C_n)$ which is composed of i runs of length 1 such that $1 \leq s_1 < s_2 < \dots < s_i \leq n$. Also let B be a j- subset of the set $\{x_{s_1+t}, x_{s_2+t}, \dots, x_{s_i+t}\} \mod n$ such that

$$\Sigma = (\Delta_t(C_n))_{\text{Vert}(\Lambda) \cup B}. \tag{5.11}$$

Suppose $\Lambda = \langle S_1, S_2, \dots, S_{\alpha+\beta} \rangle$, such that for $k = 1, 2, \dots, \alpha+\beta$, S_k is an induced subcollection of R'_k which consists of y_k runs of length one. By (5.11) we have $y_k \neq 0$ for all k. Now we prove the following claims for each $k \in \{1, 2, \dots, \alpha+\beta\}$.

a. $F_{h_k} \in \Lambda$. Suppose $1 \leq k \leq \alpha+\beta$. If $p_k = 0$ we are clearly done, so consider the case $p_k \neq 0$.

   Assume $F_{h_k} \notin \Lambda$. Since F_{h_k} is the only facet of Σ which contains x_{h_k} we can conclude $x_{h_k} \notin$ Vert(Λ). From (5.11), it follows that $x_{h_k} \in \{x_{s_1+t}, x_{s_2+t}, \dots, x_{s_i+t}\}$, so

$$x_{h_k} = x_{s_a+t} \mod n \text{ for some } a. \tag{5.12}$$

On the other hand we know

$$\begin{aligned} F_{s_a} &= \{x_{s_a}, x_{s_a+1}, \ldots, x_{s_a+t-1}\} \mod n \\ F_{s_a+1} &= \{x_{s_a+1}, x_{s_a+2}, \ldots, x_{s_a+t}\} \mod n. \end{aligned}$$

Since R'_k is an induced connected component of Σ, by (5.12) we can conclude $x_{h_k} \in F_{s_a+1}$ and $F_{s_a}, F_{s_a+1} \in R'_k$. However, we know F_{h_k} is the only facet of R'_k which contains x_{h_k} and so $F_{s_a+1} = F_{h_k}$ and then $s_a + 1 = h_k \mod n$. This and (5.12) imply that $t = 1 \mod n$, which contradicts our assumption $2 \leq t \leq n$.

b. *If $F_u \in S_k$ for some u and $F_{u+t+1} \in R'_k$, then $F_{u+t+1} \in S_k$.* Assume $F_{u+t+1} \notin S_k$ and $F_{u+t+1} \in R'_k$. Let

$$r_0 = \min\{r : r > u, F_r \in S_k \mod n\}.$$

Since S_k consists of runs of length one we can conclude $r_0 \geq u+t+1$. Since $r_0 \neq u+t+1$ we have $r_0 \geq u+t+2$. But then

$$x_{u+t+1} \notin \text{Vert}(\Lambda) \cup \{x_{s_1+t}, x_{s_2+t}, \ldots, x_{s_i+t}\}$$

and therefore $x_{u+t+1} \notin \text{Vert}(\Sigma)$ which is a contradiction.

Now for each k, by (a) we have $F_{h_k} \in \Lambda$ and from repeated applications of (b) we find that

$$F_{h_k+f(t+1)} \in S_k \quad \text{for } f = \begin{cases} 1, 2, \ldots, p_k & 1 \leq k \leq \alpha \\ 1, 2, \ldots, q_{k-\alpha} & \alpha+1 \leq k \leq \alpha+\beta. \end{cases}$$

So $R_k \subseteq S_k$. On the other hand S_k consists of runs of length one, so no other facet of R'_k can be added to it, and therefore $S_k = R_k$ for all k. We conclude that $\Lambda = \Gamma$ and we are therefore done. The last claim of the proposition is also apparent from this proof. □

We are now ready to compute the remaining Betti numbers.

Theorem 5.13. *Let n, i, j and t be integers such that $n \geq 2$, $2 \leq t \leq n$, and $ti + j < n$. Then*

i. For the cycle C_n

$$\beta_{i+j,ti+j}(R/I_t(C_n)) = \frac{n}{n-it}\binom{i}{j}\binom{n-it}{i}$$

ii. For the line graph L_n

$$\beta_{i+j,ti+j}(R/I_t(L_n)) = \binom{i}{j}\binom{n-it}{i} + \binom{i-1}{j}\binom{n-it}{i-1}$$

Proof. If $I = I_t(C_n)$ (or $I = I_t(L_n)$), from Theorem 5.3, $\beta_{i+j,ti+j}(R/I)$ is the number of $(i+j, ti+j)$-eligible subcollections of $\Delta_t(C_n)$ (or $\Delta_t(L_n)$). We consider two separate cases for C_n and for L_n.

i. For the cycle C_n, suppose $\mathcal{R}_{(i)}$ denotes the set of all induced subcollections of $\Delta_t(C_n)$ which are composed of i runs of length one. By propositions 5.11 and 5.12 there exists a one to one correspondence between the set of all $(i+j, ti+j)$-eligible subcollections of $\Delta_t(C_n)$ and the set

$$\mathcal{R}_{(i)} \times \binom{[i]}{j}$$

where $\binom{[i]}{j}$ is the set of all j-subsets of a set with i elements. By Corollary 5.4 we have $|\mathcal{R}_{(i)}| = \beta_{i,ti}$ and since $|\binom{[i]}{j}| = \binom{i}{j}$ and so we apply Corollary 5.9 to observe that

$$\beta_{i+j,ti+j}(R/I_t(C_n)) = \binom{i}{j}\beta_{i,ti}(R/I_t(C_n)) = \frac{n}{n-it}\binom{i}{j}\binom{n-it}{i}.$$

ii. For the line graph L_n, recall that

$$\Delta_t(L_n) = \langle F_1, \ldots, F_{n-t+1}\rangle.$$

Let $\Lambda = \langle F_{c_1}, F_{c_2}, \ldots, F_{c_i}\rangle$ be the induced subcollection of $\Delta_t(L_n)$ which is composed of i runs of length 1 and Vert$(\Lambda) \subset \mathcal{X} \setminus \{x_n\}$, so that it is also an induced subcollection of $\Delta_t(L_{n-1})$. Also let A be a j - subset of $\{x_{c_1+t}, x_{c_2+t}, \ldots, x_{c_i+t}\} \mod n$. So by Propositions 5.11 and 5.12 the induced subcollections on Vert$(\Lambda) \cup A$ are $(i+j, ti+j)$-eligible and if one denotes these induced subcollections by $\mathcal{B}$ we have the following bijection

$$\mathcal{B} \rightleftharpoons \binom{[i]}{j} \times \{\Gamma \subset \Delta_t(L_{n-1}) : \Gamma \quad \text{is composed of } i \text{ runs of length } 1\}. \tag{5.13}$$

We make the following claim:

Claim: *Let Γ be an $(i+j, ti+j)$-eligible subcollection of $\Delta_t(L_n)$ which contains a run $\mathcal{R}$ with $F_{n-t+1} \in \mathcal{R}$. Then $\Gamma \in \mathcal{B}$ if and only if $|\mathcal{R}| = 2 \mod t+1$.*

Proof of Claim. Let $\Gamma \in \mathcal{B}$ and assume that $\Lambda = \langle F_{c_1}, F_{c_2}, \ldots, F_{c_i}\rangle$ is the subcollection of $\Delta_t(L_{n-1})$ used to build Γ as described above. Then we must have $c_i = n-t$. Now, the run $\mathcal{R}$ contains F_{n-t+1} and F_{n-t}.

If $|\mathcal{R}| > 2$, then $c_{i-1} = n-2t-1$ and $x_{c_{i-1}+t} = x_{n-t-1} \in A$ and from Lemma 5.10 we can see that another $t+1$ facets $F_{n-2t-1}, \ldots, F_{n-t-1}$ are in $\mathcal{R}$. If we have all elements of $\mathcal{R}$, we stop, and otherwise, we continue the same way. At each stage $t+1$ new facets are added to $\mathcal{R}$ and therefore in the end $|\mathcal{R}| = 2 \mod t+1$.

Conversely, if $|\mathcal{R}| = (t+1)q+2$ then let $\Lambda = \langle F_{c_1}, F_{c_2}, \ldots, F_{c_i}\rangle$ be the unique subcollection of $\Delta_t(L_n)$ consisting of i runs of length one from which we can build Γ. Since $F_{n-t-1} \in \mathcal{R}$, we must have $c_i = n-t$ or $c_i = n-t+1$.

If $c_i = n-t$, then we are done, since Λ will be subcollection of $\Delta_t(L_{n-1})$ and so $\Gamma \in \mathcal{B}$. If $c_i = n-t+1$, then R has one facet F_{n-t+1} and if $x_{c_{i-1}+t} \in A$, then by Lemma 5.10 $\mathcal{R}$ gets an additional $t+1$ facets. And so on: for each c_u either 0 or $t+1$ facets are contributed to $\mathcal{R}$. Therefore, for some p, $|\mathcal{R}| = (t+1)p+1$ which is a contradiction. This settles our claim. ☐

We now denote the set of remaining $(i+j, ti+j)$-eligible induced subcollections of $\Delta_t(L_n)$ by $\mathcal{C}$. First we note that $\mathcal{C}$ consists of those induced subcollections which contain F_{n-t+1} and are not in $\mathcal{B}$. Also, if $j = i$, then a $(2i, (t+1)j)$-eligible subcollection Γ of $\Delta_t(L_n)$ would have no runs of length 1, as the equations in Definition 5.2 would give $\alpha = 0$. So $\Gamma \in \mathcal{C}$ and we can assume from now on that $j < i$.

We consider $\Lambda = \langle F_{c_1}, F_{c_2}, \ldots, F_{c_{i-1}}\rangle \subset \Delta_t(L_n)$ which is composed of $i-1$ runs of length 1 with Vert$(\Lambda) \subset \mathcal{X} \backslash F_{n-t+1} \cup \{x_{n-t}\}$. If A is a j-subset of the set $\{x_{c_1+t}, x_{c_2+t}, \ldots, x_{c_{i-1}+t}\}$, we claim that the induced subcollection Γ on Vert$(\Lambda) \cup A \cup F_{n-t+1}$ belongs to $\mathcal{C}$.

Suppose $\mathcal{R}$ is the run in Γ which includes F_{n-t+1}. If $|\mathcal{R}| \neq 1$ then $c_{i-1} + t = n - t$ which implies that $c_{i-1} = n-2t$. By Lemma 5.10 we see that $t+1$ facets $F_{n-2t}, F_{n-2t+1}, \ldots, F_{n-t}$ are added to $\mathcal{R}$. If these facets are not all the facets of $\mathcal{R}$ then with the same method we can see that in each step $t+1$ new facets will be added to $\mathcal{R}$ and since $F_{n-t+1} \in \mathcal{R}$ we can conclude $|\mathcal{R}| = 1 \mod t+1$. Therefore $\Gamma \notin \mathcal{B}$.

Now we only need to show that Γ is an $(i+j, ti+j)$-eligible induced subcollection. By Proposition 5.11 the induced subcollection Γ' on Vert$(\Lambda) \cup A$ is an $(i-1+j, t(i-1)+j)$-eligible induced subcollection. Suppose Γ' is composed of runs $\mathcal{R}_1, \mathcal{R}_2, \ldots, \mathcal{R}_{\alpha'+\beta'}$ and then we have

$$\begin{cases} t(i-1)+j & = (t+1)(P'+Q') + t(\alpha'+\beta') + \beta' \\ i-1+j & = 2(P'+Q') + 2\beta' + \alpha' \end{cases} \Longrightarrow \begin{cases} i-1 & = P'+Q'+\alpha'+\beta' \\ j & = P'+Q'+\beta' \end{cases} \tag{5.14}$$

So Γ consists of all or all but one of the runs $\mathcal{R}_1, \mathcal{R}_2 \ldots, \mathcal{R}_{\alpha'+\beta'}$ as well as $\mathcal{R}$ where $\mathcal{R}$ is the run which includes F_{n-t+1}.

As we have seen $|\mathcal{R}| = 1 \mod t+1$. If we suppose $|\mathcal{R}| = 1$ then we can claim that Γ is composed of $\alpha = \alpha'+1$ runs of length 1 and $\beta = \beta'$ runs of length $2 \mod t+1$, and with $P = P'$ and $Q = Q'$, by (5.14) we have Γ is an $(i+j, ti+j)$-eligible induced subcollection. Now assume $|\mathcal{R}| = (t+1)p+1$, so clearly we have $F_{n-2t} \in \Lambda$ and $x_{n-t} \in A$. Let $\mathcal{R}'$ be the induced subcollection on Vert$(\mathcal{R}) \setminus F_{n-t+1}$. Then clearly we have $\mathcal{R}'$ is a run in Γ' and since the only facets which belong to $\mathcal{R}$ but not to $\mathcal{R}'$ are the t facets $F_{n-2t+2}, \ldots, F_{n-t+1}$ we have

$$|\mathcal{R}'| = (t+1)p + 1 - t = (t+1)(p-1) + 2 \tag{5.15}$$

Therefore we have shown the run in Γ which includes F_{n-t+1} has been generated by a run of length $2 \mod (t+1)$ in Γ'. Using (5.14) we can conclude Γ consists of $\alpha = \alpha'+1$ runs of length 1 and $\beta = \beta'-1$ runs of length $2 \mod t+1$. We set $P = P'+p$ and $Q = Q'-(p-1)$, and use (5.15) to conclude that

$$\begin{cases} P+Q+\alpha+\beta = (P'+p) + (Q'-p+1) + (\alpha'+1) + (\beta'-1) = i \\ P+Q+\beta = (P'+p) + (Q'-p+1) + (\beta'-1) = j. \end{cases}$$

Therefore $\Gamma \in \mathcal{C}$ as we had claimed.

Conversely, let $\Gamma \in \mathcal{C}$ then one can consider the induced subcollection Γ' on Vert$(\Gamma) \setminus F_{n-t+1}$. Assume Γ is composed of runs $\mathcal{R}_1, \mathcal{R}_2 \ldots, \mathcal{R}_{\alpha+\beta}$, so that $\mod t+1$, $\mathcal{R}_h$ is a run of length 1 if $h \leq \alpha$ and length 2 otherwise.

Suppose $\mathcal{R}_h$ is the run which includes F_{n-t+1}. By our assumption we have $|\mathcal{R}_h| = 1 \mod t+1$, so $h \leq \alpha$. If $|\mathcal{R}_h| = 1$ then $\mathcal{R}_h \notin \Gamma'$ and therefore we delete one run of length one from Γ to obtain Γ', in which case Γ' is $(i-1+j, t(i-1)+j)$-eligible.

If $|\mathcal{R}_h| = (t+1)p_h + 1 > 1$ then the t facets $F_{n-2t+2}, \ldots, F_{n-t+1} \in \mathcal{R}_h$ do not belong to Γ'. So Γ' consists of $\alpha+\beta$ runs $\mathcal{R}_1, \ldots, \widehat{\mathcal{R}_h}, \ldots, \mathcal{R}_{\alpha+\beta}, \mathcal{R}'_h$ where

$$|\mathcal{R}'_h| = (t+1)p_h + 1 - t = (t+1)(p_h-1) + 2.$$

Setting $\alpha' = \alpha-1$, $\beta' = \beta+1$, $P' = P - p_h$ and $Q' = Q + p_h - 1$ it follows that Γ' is $(i-1+j, t(i-1)+j)$-eligible. By Proposition 5.12 there exists a unique induced subcollection $\Lambda = \langle F_{c_1}, F_{c_2}, \ldots, F_{c_{i-1}} \rangle$ of $\Delta_t(L_{n-t-1})$ which is composed of $i-1$ runs of length one and a j subset A of $\{x_{c_1+t}, \dots, x_{c_{i-1}+t}\}$ such that Γ' equals to induced subcollection on $\mathrm{Vert}(\Lambda) \cup A$. So Γ is the induced subcollection on $\mathrm{Vert}(\Lambda) \cup A \cup F_{n-t+1}$. Therefore there is a one to one correspondence between elements of $\mathcal{C}$ and

$$\binom{[i-1]}{j} \times \{\Gamma \subset \Delta_t(L_{n-t-1}) : \Gamma \quad \text{is composed of } i-1 \text{ runs of length } 1\} \tag{5.16}$$

By (5.13), (5.16) and Corollary 5.9 (iii) we have

$$\begin{aligned} \beta_{i+j,ti+j}(R/I) &= |\mathcal{B}| + |\mathcal{C}| \\ &= \tbinom{i}{j}\beta_{i,ti}(R/I_t(L_{n-1})) + \tbinom{i-1}{j}\beta_{i-1,t(i-1)}(R/I_t(L_{n-t-1}) \\ &= \tbinom{i}{j}\tbinom{n-it}{i} + \tbinom{i-1}{j}\tbinom{n-it}{i-1}. \end{aligned}$$

□

Finally, we put together Theorem 5.3, Proposition 5.5, Theorem 5.1, (5.1) and Theorem 2.8. Note that the case $t = 2$ is the case of graphs which appears in Jacques [8].

Theorem 5.14 (Betti numbers of path ideals of lines and cycles). *Let n, t, p and d be integers such that $n \geq 2$, $2 \leq t \leq n$, $n = (t+1)p + d$, where $p \geq 0$, $0 \leq d \leq t$. Then*

i. The $\mathbb{N}$-graded Betti numbers of the path ideal of the graph cycle C_n are given by

$$\beta_{i,j}(R/I_t(C_n)) = \begin{cases} t & j = n,\ d = 0,\ i = 2\left(\dfrac{n}{t+1}\right) \\[2ex] 1 & j = n,\ d \neq 0,\ i = 2\left(\dfrac{n-d}{t+1}\right) + 1 \\[2ex] \dfrac{n}{n - t\left(\frac{j-i}{t-1}\right)} \dbinom{\frac{j-i}{t-1}}{\frac{ti-j}{t-1}} \dbinom{n - t\left(\frac{j-i}{t-1}\right)}{\frac{j-i}{t-1}} & \begin{cases} j < n,\ i \leq j \leq ti, \text{ and} \\ 2p \geq \dfrac{2(j-i)}{t-1} \geq i \end{cases} \\[2ex] 0 & \text{otherwise.} \end{cases}$$

ii. The $\mathbb{N}$-graded Betti numbers of the path ideal of the line graph L_n are nonzero and equal to

$$\beta_{i,j}(R/I_t(L_n)) = \dbinom{\frac{j-i}{t-1}}{\frac{ti-j}{t-1}} \dbinom{n - t\left(\frac{j-i}{t-1}\right)}{\frac{j-i}{t-1}} + \dbinom{\frac{j-i}{t-1} - 1}{\frac{ti-j}{t-1}} \dbinom{n - t\left(\frac{j-i}{t-1}\right)}{\frac{j-i}{t-1} - 1}$$

if and only if

(a) $j \leq n$ and $i \leq j \leq ti$;

(b) If $d < t$ then $p \geq \dfrac{j-i}{t-1} \geq i/2$ where both inequalities cannot be $=$ at the same time;

(c) If $d = t$ then $(p+1) \geq \dfrac{j-i}{t-1} \geq i/2$ where both inequalities cannot be $=$ at the same time.

Proof. We only need to make sure we have the correct conditions for the Betti numbers to be nonzero.

i. When $j < n$, $\beta_{i,j}(R/I_t(C_n)) \neq 0 \iff$

$$
\begin{aligned}
&\iff \begin{cases} \dfrac{j-i}{t-1} \geq \dfrac{ti-j}{t-1} \\[2ex] n - \dfrac{t(j-i)}{t-1} \geq \dfrac{j-i}{t-1} \end{cases} \\[2ex]
&\iff \begin{cases} 2j \geq (t+1)i \\[1ex] n \geq \left(\dfrac{t+1}{t-1}\right)(j-i) \end{cases} \\[2ex]
&\iff \begin{cases} 2j \geq (t+1)i \\[1ex] (t+1)p + d \geq \left(\dfrac{t+1}{t-1}\right)(j-i) \end{cases} \qquad (5.17) \\[2ex]
&\iff \begin{cases} 2j \geq (t+1)i \\[1ex] p + \dfrac{d}{t+1} \geq \dfrac{j-i}{t-1} \end{cases} \\[2ex]
&\iff 2p \geq \frac{2(j-i)}{t-1} \geq i \qquad \text{as } d < t+1
\end{aligned}
$$

ii. $\beta_{i,j}(R/I_t(L_n)) \neq 0 \iff$

$$
\begin{aligned}
&\iff \begin{cases} \dfrac{j-i}{t-1} \geq \dfrac{ti-j}{t-1} \\[2ex] n - \dfrac{t(j-i)}{t-1} \geq \dfrac{j-i}{t-1} - 1 \end{cases} \qquad \text{both } \geq \text{ cannot be } = \text{ at the same time} \\[2ex]
&\iff \begin{cases} 2j \geq (t+1)i \\[1ex] n + 1 \geq \left(\dfrac{t+1}{t-1}\right)(j-i) \end{cases} \qquad \text{both } \geq \text{ cannot be } = \text{ at the same time} \\[2ex]
&\iff \begin{cases} 2j \geq (t+1)i \\[1ex] (t+1)p + d + 1 \geq \left(\dfrac{t+1}{t-1}\right)(j-i) \end{cases} \qquad \text{both } \geq \text{ cannot be } = \text{ at the same time} \qquad (5.18) \\[2ex]
&\iff \begin{cases} 2j \geq (t+1)i \\[1ex] p + \dfrac{d+1}{t+1} \geq \dfrac{j-i}{t-1} \end{cases} \qquad \text{both } \geq \text{ cannot be } = \text{ at the same time} \\[2ex]
&\iff \begin{cases} 2p \geq \dfrac{2(j-i)}{t-1} \geq i & \text{if } d < t \\[2ex] 2(p+1) \geq \dfrac{2(j-i)}{t-1} \geq i & \text{if } d = t \end{cases} \qquad \text{both } \geq \text{ cannot be } = \text{ at the same time in each line.}
\end{aligned}
$$

□

We end the paper with the computation of the projective dimension and regularity of path ideals of lines and cycles. The projective dimension of lines (Part i below) was computed by He and Van Tuyl in [7] using different methods. The case $t = 2$ is the case of graphs which appears in Jacques [8].

Part ii of the following Corollary reproves Theorem 5.3 in [1] which computes the Castelnuovo-Mumford regularity of path ideal of a line.

Corollary 5.15 (Projective dimension and regularity of path ideals of lines and cycles). *Let n, t, p and d be integers such that $n \geq 2$, $2 \leq t \leq n$, $n = (t+1)p + d$, where $p \geq 0$, $0 \leq d \leq t$. Then*

i. The projective dimension of the path ideal of a graph cycle C_n or line L_n is given by

$$pd(R/I_t(C_n)) = \begin{cases} 2p+1 & d \neq 0 \\ 2p & d = 0 \end{cases} \qquad pd(R/I_t(L_n)) = \begin{cases} 2p & d \neq t \\ 2p+1 & d = t \end{cases}$$

ii. The regularity of the path ideal of the graph cycle C_n or line L_n is given by

$$reg(R/I_t(C_n)) = (t-1)p + d. \qquad reg(R/I_t(L_n)) = \begin{cases} p(t-1) & d < t \\ (p+1)(t-1) & d = t \end{cases}$$

Proof. i. From Hochster's formula (Theorem 2.8) we know the maximum degree for the Betti numbers is $j = n$ and that can only happen at the far end of the resolution. Therefore, in the case of the cycle C_n, the claimed formula for projective dimension follows from Theorem 5.14. For the case L_n for $d = t$ by applying Theorem 5.14 we have

$$\beta_{2p+1,n}(R/I_t(L_n)) = \binom{p+1}{p}\binom{p}{p+1} + \binom{p}{p}\binom{p}{p} = 1 \neq 0.$$

Then by same argument as above we can conclude that $pd(R/I_t(L_n)) = 2p+1$.

Now we suppose $d \neq t$. From (5.18) we can see that if $\beta_{i,j}(R/I_t(L_n)) \neq 0$ then $2p \geq i$ and therefore $pd(R/I_t(L_n)) \leq 2p$. On the other hand, by applying Theorem 5.14 again, we can see that

$$\beta_{2p,p(t+1)}(R/I_t(L_n)) = \binom{p}{p}\binom{p+d}{p} + \binom{p-1}{p}\binom{p}{p} = \binom{p+d}{p} \neq 0.$$

Therefore $pd(R/I_t(L_n)) \geq 2p$ and we have $pd(R/I_t(L_n)) = 2p$.

ii. By definition, the regularity of a module M is $\max\{j - i \mid \beta_{i,j}(M) \neq 0\}$. By Theorem 5.14, we know exactly when the graded Betti numbers of $R/I_t(L_n)$ and $R/I_t(C_n)$ are nonzero.

(a) In the case of C_n if $d = 0$ then

$$reg(R/I_t(C_n)) = \max\{n - \frac{2n}{t+1}, (t-1)p\} = \max\{(t+1)p - 2p, (t-1)p\} = (t-1)p$$

and if $d \neq 0$ then

$$reg(R/I_t(C_n)) = \max\{n - \frac{2(n-d)}{t+1} - 1, (t-1)p\} = \max\{(t+1)p + d - 2p, (t-1)p\} = (t-1)p + d.$$

The formula now follows.

(b) In the case of L_n, the formula follows directly from (5.18).

□

References

[1] R. Bouchat, H. T. Ha, and A. O' Keefe. Path ideals of rooted trees and their graded betti numbers. *math.AC math.CO*, page 18, 2010.

[2] W. Bruns and J. Herzog. *Cohen-Macaulay rings*. Cambridge University Press, 1993.

[3] A. Conca and E. De Negri. M-sequences, graph ideals and ladder ideals of linear types. *J. Algebra*, 211:599–624, 1999.

[4] D. R. Grayson and M. E. Stillman. Macaulay 2, a software system for research in algebraic geometry. Technical report, http://www.math.uiuc.edu/Macaulay2/.

[5] R. P. Grimaldi. *Discrete and Combinatorial Mathematics, An applied Introduction*. 2003.

[6] A. Hatcher. *Algebraic Topology*. Cambridge University Press, 2002.

[7] J. He and A. Van Tuyl. Algebraic properties of path ideal of a tree. *Comm. Algebra*, 38:1725–1742, 2010.

[8] S. Jacques. Betti numbers of graph ideals. *PhD thesis available on line 'http://arxive.org/math/0410107*, 2004.

[9] E. Miller and B. Sturmfels. *Combinatorial Commutative Algebra*. Spring -Verlag, New York, 2005.

[10] R. P. Stanley. *Enumerative Combinatorics*. Book manuscript, 1994.

[11] CoCoA Team. Cocoa: a system for doing computations in commutative algebra. Technical report, Available at http://cocoa.dima.unige.it.

[12] R. H. Villarreal. *Monomial Algebras*. Marcel Dekker Inc, 2001.